



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|--|---|
| (51) International Patent Classification ⁶ : A61K 38/28, 31/19, 31/35 | A1 | (11) International Publication Number: WO 00/16798 (43) International Publication Date: 30 March 2000 (30.03.00) |
| (21) International Application Number: PCT/AU99/00786 (22) International Filing Date: 17 September 1999 (17.09.99) (30) Priority Data: PP 6091 22 September 1998 (22.09.98) AU (71) Applicant (for all designated States except US): CURTIN UNIVERSITY OF TECHNOLOGY [AU/AU]; Kent Street, Bentley, W.A. 6102 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only): HELMERHORST, Erik [AU/AU]; 5 Selsted Place, Glendalough, W.A. 6016 (AU). PLEWRIGHT, Brian, Scott [AU/AU]; 253 Grand Promenade, Doubleview, W.A. 6018 (AU). (74) Agent: WRAY & ASSOCIATES; 239 Adelaide Terrace, Perth, W.A. 6000 (AU). | (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. | |

(54) Title: USE OF NON-PEPTIDYL COMPOUNDS FOR THE TREATMENT OF INSULIN RELATED AILMENTS

(57) Abstract

The present invention relates to the use of at least a non-peptidyl compound as a biological modulator of insulin activity or insulin-related activity, which compound possesses ionic and hydrophobic chemical moieties spatially located so as to mimic at least an ionic and hydrophobic amino acid residue of insulin, which amino acids are associated with the binding of insulin to its receptor.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | KR | Republic of Korea | PL | Poland | | |
| CN | China | KZ | Kazakhstan | PT | Portugal | | |
| CU | Cuba | LC | Saint Lucia | RO | Romania | | |
| CZ | Czech Republic | LI | Liechtenstein | RU | Russian Federation | | |
| DE | Germany | LK | Sri Lanka | SD | Sudan | | |
| DK | Denmark | LR | Liberia | SE | Sweden | | |
| EE | Estonia | | | SG | Singapore | | |

Use of Non-Peptidyl Compounds for the Treatment of
Insulin Related Ailments

Field of the Invention

The present invention relates to the use of one or more non-peptidyl compounds
5 for the treatment of a patient suffering from one or more insulin related ailments.

Background Art

General

Those skilled in the art will appreciate that the invention described herein is
susceptible to variations and modifications other than those specifically described.
10 It is to be understood that the invention includes all such variation and
modifications. The invention also includes all of the steps, features, compositions
and compounds referred to or indicated in the specification, individually or
collectively, and any and all combinations or any two or more of the steps or
features.

15 The present invention is not to be limited in scope by the specific embodiments
described herein, which are intended for the purpose of exemplification only.
Functionally equivalent products, compositions and methods are clearly within the
scope of the invention as described herein.

Bibliographic details of the publications numerically referred to in this specification
20 are collected at the end of the description. All references cited, including patents
or patent applications are hereby incorporated by reference. No admission is
made that any of the references constitute prior art.

As used herein the term "derived from" shall be taken to indicate that a specific
integer may be obtained from a particular source *albeit* not necessarily directly
25 from that source.

Throughout this specification and the claims which follow, unless the context
requires otherwise, the word "comprise", or variations such as "comprises" or
"comprising", will be understood to imply the inclusion of a stated integer or group
of integers but not the exclusion of any other integer or group of integers.

Scientific

It is generally accepted that agents such as hormones and growth factors elicit their biological functions by binding to specific recognition sites, or receptors, in the plasma membranes of their target cells. This typically causes a
5 conformational change in the receptor, and triggers secondary cellular responses which may result in the activation or inhibition of intracellular processes. Such agents are often referred to as biological modifiers. They may be categorised into two classes according to their activity. Agents which have a stimulatory activity are termed agonists, and those which inhibit the effect of the original ligand are
10 termed antagonists.

The discovery of biological modifiers that differ in structure from the original ligand may be medically useful. Such compounds may have slightly different spectra of biological activity, allowing them to be used in very specific situations. Compounds that are chemically simpler than the agent which they mimic are
15 potentially produced in greater quantities and at lower costs. Compounds that are more chemically robust than the original ligand may be administered by more convenient means. For example, peptidyl ligands cannot be taken orally as they will be broken down in the digestive tract, whereas a non-peptidyl agonist may potentially be administered by this route.

20 Insulin is a peptidyl ligand, and the identification of agonists and antagonists thereof has important pharmaceutical applications in the treatment of insulin related ailments.

Insulin is regarded as the most important regulatory hormone involved in maintaining glucose homeostasis. *In vivo*, insulin is produced in the pancreatic β -
25 cells of the Islets of Langerhans. It is secreted from these cells in response to a wide range of nutrients, hormones and neurotransmitters. However, glucose is considered the most important regulator of insulin release. Following its release insulin is carried systemically to target tissues such as liver, muscle and fat. It promotes the uptake of glucose in the peripheral tissues and inhibits
30 gluconeogenesis in the liver. Insulin also is involved in: promoting the synthesis of

glycogen, lipid, and protein; gene transcription and mRNA turnover, and the transport of specific amino acids and ions.

Insulin works by first binding to the insulin receptor on the cell surface of its target tissues. The insulin receptor is a specific transmembrane glycoprotein composed of two α -subunits and two β -subunits linked by disulfide bonds. The α -subunits are located extracellularly and contain the insulin binding domain. The β -subunits of the receptor pass through the plasma membrane and have an intrinsic tyrosine kinase activity associated with their intracellular domain. The X-ray coordinates of part of the tyrosine kinase domain has been resolved (1, 2), however, the coordinates detailing the structure of the α -subunits are not yet available. Some inferences about the structure of some domains of the α -subunit can be made from fibronectin structures of homologous protein domains resolved by NMR spectroscopy and from the X-ray structure of the closely related IGF-1 receptor (3). Most recently, the general quaternary structure of the insulin receptor has been resolved using electron cryomicroscopy (4). These studies indicate that one insulin molecule probably binds and effectively crosslinks a L1-cysteine rich domain of one α -subunit and the L2 domain of the other α -subunit. This viewpoint is supported by models derived from the complex kinetics of insulin binding and the models of receptor oligomerisation described below. Following insulin binding to the α -subunit, the β -subunit is autophosphorylated on specific tyrosine residues and this promotes the tyrosine kinase activity of the receptor.


The interaction of insulin with its receptor leading to receptor activation is a complex, key initial event in insulin action. The kinetics of insulin binding to its receptor have been discussed in a plethora of literature over the past twenty years and many different models have been proposed to explain the interaction. Much debate still exists over which of these proposed models effectively describes the correct binding kinetics.

The simplest model proposes that the insulin receptor population is characterised by a single class of homogeneous, non-interacting binding sites and that insulin action is directly proportional to the fraction of these receptors that are occupied. Other studies confirm a single class of receptors, with reported dissociation

constants in the order of 0.5 to 5 nM, depending on the source of insulin receptors and experimental conditions employed (eg. temperature).

- An alternate model is based on studies that report curvilinear Scatchard plots and assume multiple classes of binding sites. R. Kahn *et al.* (5) propose two specific
5 receptor sites; a high affinity-low capacity site and a low affinity-high capacity site. However, in a recent review, B.J. Hammond *et al.* (6), question the validity of this and suggest that because these two binding sites exist on each receptor, there is no logical reason why they would possess different affinities or more importantly widely differing concentrations.
- 10 Another model explaining the non-linear nature of Scatchard plots assumes negative cooperativity. P. De Meyts *et al.* (7), suggest that the filling of empty receptor sites by unlabelled insulin increases the dissociation of labelled insulin from other sites, leading to curvilinear Scatchard plots. This was interpreted as resulting from site-site interactions. However, E. Helmerhorst (8) demonstrates
15 that insulin enhances its own dissociation even when binding is characterised by linear Scatchard plots. Additionally, R. Pollet *et al.* (9) recognise that the dissociation rate of insulin is nearly independent of receptor occupancy, with enhanced dissociation of bound ¹²⁵I insulin by native insulin occurring in certain conditions where binding occupancy is decreased. D. Donner (10) also question
20 the theory of negative cooperativity because correction for insulin degradation products result in linear Scatchard plots, suggesting that heterogeneous distribution of binding may in fact be due to insulin degradation rather than negative cooperative interactions.

- Another two-state model, proposed by R. Corin and D. Donner (11) explains the
25 kinetics observed upon insulin/receptor dissociation and is illustrated below:



The hormone (H) interacts with the receptor (R_1) to result in the formation of a complex (HR_1), with subsequent rapid dissociation (k_{-1}). Stabilisation of the complex (HR_1) is through conversion (k_2) to another state, from which hormone dissociation is slow (k_{-2}). R. Corin and D. Donner (11) postulate that increasing occupancy time favours the formation of the second complex at the expense of the first. Thus, this model supports the kinetic data that identifies a constant association rate, but a dissociation rate which decreases with time. As a result of a decreased dissociation rate, the dissociation constant is also reduced, therefore providing an explanation for the presence of receptors of two apparent different affinities. It is not known if the bound complex HR_2 may dissociate directly to H and R_2 through k_{-3} and to R_1 through k_{-4} .

A study by E. Helmerhorst and C. Yip (12) support this two-state model, with the binding of insulin to rat liver membranes producing Scatchard plots of both a linear and curvilinear nature. Insulin binding to rat liver membranes at temperatures of 4 to 15°C is characterised by linear Scatchard plots, while higher temperatures result in plots that are distinctly curvilinear. E. Helmerhorst and C. Yip (12) implicate a single class of homogeneous, non-interacting binding sites of low affinity below 15°C. However, at physiological temperatures, two receptor states exist. In reference to the model of R. Corin and D. Donner (11), this suggests that insulin receptors exist primarily in one state at temperatures below 15 degrees, where $R_1 > R_2$ and k_4 is less than k_{-4} . However, as the temperature increases, these values become more similar, leading to increased R_2 , and a curvilinear Scatchard plot.

Several models are based on the original suggestion of D.B. Donner and K. Yonkers (13), that insulin binding induces a conformational change in its receptor, with a conversion to a more slowly dissociating complex. P. De Meyts (14) and L. Schaffer (15) models are consistent with this notion where the initial binding of insulin to one α -subunit of the insulin receptor is followed by the cross-linking of insulin to a second α -subunit.

Briefly, the model proposes that insulin contains two binding surfaces on each α -subunit which can interact independently with corresponding surfaces on another α -subunit of the receptor. These binding sites, termed $\alpha 1$ and $\alpha 2$, have differing affinities with $\alpha 1$ having a higher affinity than the $\alpha 2$ region. If insulin binds to the $\alpha 1$ site on one α -subunit, it is suggested that this induces a conformational change in the receptor which ultimately allows the insulin molecule to interact with the $\alpha 2$ site on the second subunit. Consequently, the two α -subunits are cross-linked (homo dimerisation), resulting in high affinity binding between insulin and its receptor and subsequent activation of the receptor. However, although another insulin molecule may potentially bind to the $\alpha 1$ site that is vacant, the previous conformational change prevents it from interacting with the second $\alpha 2$ site, causing this interaction to be one of lower affinity. L. Schaffer (15) suggests that negative interactions between the cross-linked tracer insulin and a native insulin molecule bound at this second $\alpha 1$ site may explain the accelerated dissociation in the presence of high concentrations of insulin. However, B.J. Hammond *et al.* (6) discount this explanation and propose that the acceleration is in fact due to the destabilisation of the cross-linked formation upon the second insulin molecule binding. At very high concentrations of unlabelled insulin, acceleration of tracer insulin dissociation from receptor is inhibited and is attributed by both P. De Meyts (14) and L. Schaffer (15) to monovalent insulin binding of a third insulin molecule to the vacant low affinity $\alpha 2$ site (B.J. Hammond *et al.* (6)). Therefore, although the stoichiometry of insulin binding to its receptor can support up to three molecules of insulin bound per $\alpha 2 \beta 2$ receptor, the more complex 1:2 and 1:3 complexes occur at supra physiological concentrations of insulin and may not be physiologically relevant.

Another factor that may contribute to the complexity of the interaction between insulin and its receptor is the possible involvement of other non-receptor related molecules. J.T. Harmon *et al.* (16), propose the presence of a membrane protein affinity regulator of insulin binding in rat liver membranes from ob/ob mice. They report that the effect of this protein is to cause a lowering of insulin affinity for its receptor. In the presence of high salt, high pH or purification of receptors on lectin columns, insulin's apparent affinity for its receptor increases. This effect is reversed by dilution of salt or reduction of pH. A similar effect in human placental plasma membranes has been observed and the apparent increase in affinity that has been determined is due to a two-fold increase in the high affinity component of insulin binding rather than an increase in affinity of either the high or low affinity site. R.A. Kohanski and M.D. Lane (17) also demonstrate the presence of a peripheral membrane glycoprotein which is involved in the modulation of insulin binding, while J. Maturo and M. Hollenberg (18) observe a reduction in binding affinity when purified insulin receptors are supplemented with the glycoprotein eluant from the affinity column used to purify them (also observed in our laboratory). E.R. Mortensen *et al* (19) demonstrate that mild reduction of adipocyte plasma membranes increases the proportion of high affinity insulin binding which is decreased in the presence of GTP, and is temperature and concentration dependent. Thus, the temperature effect observed by E. Helmerhorst and C. Yip (12) may be attributed to the presence of a G-protein that modulates insulin binding. More recently, the presence of a 66 kDa GTP-binding protein in human placental plasma membrane preparations has been shown to influence the binding kinetics of insulin (20). A decrease in insulin binding to the insulin receptor occurs upon pre-incubation of the insulin receptor-G-protein fraction with GTP γ S (a stable analogue of GTP), suggesting an important interaction between the insulin receptor and a G-protein. The presence of such a peripheral membrane protein has not been widely espoused in the literature, however, it may well explain some of the complexities of the binding interaction.

Receptor oligomerisation is a ubiquitous phenomenon among growth factor receptors. It may be induced by monomeric ligands such as EGF to induce conformational changes that result in receptor-receptor interactions, or bivalent ligands (eg human growth hormone (hGH)) that mediate dimerisation of

neighbouring receptors. The structure of subclass II receptors eg. insulin and IGF-1, are different because they exist as disulfide linked pairs of dimers in a heterotetrameric structure. In these cases, ligand binding to the receptor induces allosteric interaction between the two $\alpha\beta$ halves to activate the receptors.

- 5 Recently, through high resolution structural and functional studies the fundamental mechanics of ligand-induced receptor activation is beginning to be elucidated. These studies indicate that oligomerisation of receptor subunits may be all that is required for receptor activation. Moreover, the proposed models explain some of the complex kinetics observed in ligand/receptor binding and
- 10 dose-response studies of these molecules. For example, a model of homodimerisation of growth hormone receptor explains the unusual bell-shaped dose response curve which is caused by monovalent binding of hGH to each receptor monomer and subsequent prevention of dimerisation.

It is generally accepted that insulin binding induces a conformational change in

15 the insulin receptor that is responsible for receptor activation. A number of studies indicate that this conformational change may occur by a cross-linking mechanism, similar to that described above. Firstly, insulin receptors are activated by a number of anti insulin receptor antibodies, but not by monovalent Fab' fragments, which implies a receptor cross-linking mechanism of activation

20 similar to EGF receptors. Secondly, the homobifunctional cross-linking agent disuccinimide suberate forms more α_2 species in the presence than in the absence of insulin, indicating that insulin draws the α -subunits into close proximity. Thirdly, reduced $\alpha\beta$ dimer half receptors, when immobilised on wheat germ agglutinin Sepharose do not phosphorylate in the presence of insulin.

25 However, when they are free to associate in solution, insulin causes phosphorylation of the β -subunits, indicating that cross-linking of receptor $\alpha\beta$ halves must occur for receptor activation. Finally, concanavalin A induces insulin receptor activation by cross-linking receptors in intact cells, but when monovalent, only slightly induces receptor activation.

- 30 Although the dimeric insulin receptor can bind two (or more) insulin molecules (compared with growth hormone which cross-links two hGH receptors) the first

insulin molecule binds more tightly than the second molecule and maybe all that is physiologically relevant for receptor activation. In this respect, insulin receptor activation may well be synonymous with hGH activation of its receptor (or similar cytokine/hormone receptor activation). Indeed, the increased rate of dissociation
5 of tracer insulin in the presence of unlabelled insulin has been explained by analogy to the self-antagonism seen for hGH and its receptors.

Insulin binding to the two receptor α -subunits causes activation of the receptor β -subunit tyrosine kinase activity which leads to the phosphorylation of various substrates of insulin action. This includes the phosphorylation of a family of
10 insulin receptor substrates (IRS) that are believed to be the immediate downstream effector molecules of much of insulin action. At present four members of the IRS family have been discovered. It is believed that these molecules have the potential to interact with, and thereby activate, other downstream signalling molecules, leading to many of the actions of insulin.
15 However, the signalling pathways regulated by the individual substrates may vary.

Given the importance of insulin in maintaining glucose homeostasis then, it is clear that any change in the levels of insulin secreted or the responsiveness of cells to insulin may have significant consequences. Decreased secretion of insulin, or decreased responsiveness of cells to insulin results in Diabetes
20 Mellitus, a group of individual conditions, distinguished by the varying causes of hyperglycemia.

Diabetes is a complex disease with many causative factors. Hyperglycemia is a major characteristic of the disease and over time, especially if poorly controlled, leads to many complications of the disease. These complications include
25 microvascular and macrovascular diseases, retinopathy, neuropathy, stroke, hypertension, heart and kidney disease. Careful control of blood glucose levels is, therefore, a key strategy in treatment of diabetes.

Type 1 diabetes mellitus, or insulin-dependent diabetes mellitus (IDDM) is the more severe form of the disease, and is usually detected before 40 years of age.
30 IDDM results from an autoimmune disorder leading to an inability of the body to produce the insulin needed to help maintain blood glucose levels within a normal

range (the roles of insulin and how it works is described in more detail below). The symptoms of IDDM include polyuria, polyphagia, polydipsia, weight loss and drowsiness.

Presently, IDDM patients are absolutely dependent on regular injections of insulin
5 for survival. Over 20 million people worldwide are dependent upon insulin in this manner.

Several different types of human insulin are commercially available for diabetics, ranging from the fast-acting Humulin™BR and Novolin™ to slower acting treatments, such as Protamine-zinc-insulin (PZI), Neutral protamine Hagedorn
10 (NPH) insulin and Lente insulin. Insulin analogues like Humalog (LysPro) with altered properties are also available. Each of these insulin therapies have their advantages and disadvantages and so the choice of insulin therapy should be made by the patient and physician with all information about the patient's lifestyle, physical performance, and drug preferences.

15 The more common form of diabetes, representing in excess of 90% of all diagnosed cases, is referred to as Type 2 diabetes mellitus or non-insulin dependent diabetes mellitus (NIDDM). NIDDM can be triggered by both genetic and environmental factors and is often found in obese individuals. There are at least two fundamental defects associated with NIDDM. One is an increase in the
20 resistance of cells in peripheral tissue to the presence of insulin. Another is decreased secretion of insulin by the β -cells damaged by long-term, elevated blood glucose levels.

The disruption of the insulin receptor substrate (IRS) signalling system involved in mediating the cellular response to insulin also may play a role in the development
25 of NIDDM. Unlike IDDM, the clinical symptoms of NIDDM are often mild, and the condition may even be asymptomatic. NIDDM patients usually do not depend upon insulin injections for their survival. About half adequately control blood glucose levels through dietary therapy and exercise regimes. The others use various oral hypoglycemia agents such as sulfonylureas, biguanides, and α -
30 glucosidase inhibitors, or insulin, or various combinations of them to help regulate their blood glucose levels. The exact drug prescribed to a patient depends not

only on the patient's clinical characteristics, but also the pharmacological properties of the treatment.

The sulfonylureas are the most common oral hypoglycemics and are traditionally used in the treatment of non-obese sufferers of NIDDM. They promote the beta-
5 cells in the Islets of Langerhans of the pancreas to secrete insulin and so they effectively augment glucose-induced insulin secretion.

However, these drugs appear to be responsible for inducing hypoglycemic episodes in patients with the incidence of this apparently increasing when sulfonylureas are used in conjunction with alcohol, drugs which potentiate
10 sulfonylurea action, poor food intake or renal impairment. A weight gain is often associated with sulfonylurea use making these compounds undesirable for the treatment of NIDDM in overweight patients.

The biguanide class of oral hypoglycemic agents increase insulin sensitivity and therefore can be used to lower blood glucose levels in NIDDM. Their mechanism
15 of action is unclear. Phenformin is no longer used to treat NIDDM as it can cause fatal lactic acidosis. Metformin now is the only biguanide in clinical use worldwide. It is most commonly used when dietary therapy is unsuccessfully used to regulate blood glucose levels in obese patients. Metformin may be used in conjunction with sulfonylureas in instances where sulfonylurea therapy alone is inadequate or
20 it may be used in combination with insulin in the treatment of IDDM. However, the adverse side-effects of metformin therapy may include lactic acidosis, nausea, bloating, diarrhoea and abdominal cramping.

α -Glucosidase inhibitors are also used to treat NIDDM as an adjunct to dietary measures or sulfonylureas therapy. These compounds allow carbohydrate in the
25 gut to be processed more effectively by slowing down their absorption from the intestinal tract. Adverse side effects of these compounds include flatulence, diarrhoea and abdominal pain.

The thiazolidinediones are another class of compounds that may ameliorate symptoms of NIDDM. These compounds work by reducing insulin resistance at
30 the sites of insulin action in the muscle and liver. They may be used in

combination with insulin or sulfonylurea drugs but are not recommended for the treatment of IDDM. Severe adverse side-effects are rarely observed, however, some NIDDM patients fail to respond to this treatment.

Conversely, a number of clinical conditions are characterised by hyperinsulinism
5 that leads to hypoglycaemia.

Insulin or hypoglycaemic drug overdose are clinical conditions that are often difficult to manage and may require hospitalisation and several days of intensive care. Non-accidental overdose or suicide attempts are fairly rare but often lead to death or profound neurologic impairment. The key symptoms of insulin overdose
10 are hypoglycaemia, hypokalaemia and acid-base imbalance. Sulfonylurea drug overdose predominantly causes hypoglycaemia.

Insulinomas account for about 90% of all pancreatic endocrine tumours. They occur with an incidence of about 0.5 per million population and people of all ages can be affected. Early diagnosis and treatment of insulinomas is essential
15 because of their variable manifestations and potential lethality. These tumours are usually benign but synthesise and secrete insulin autonomously causing spontaneous hypoglycaemia. Symptoms may include deep coma, epilepsy, dizziness, weakness, hunger and epigastric pain.

Congenital hyperinsulinism is the most common cause of severe, persistent
20 hypoglycaemia in infants. It may be familial as up to 20% of affected families have more than one affected child. A defect in beta-cell function is the most likely explanation for the hyperinsulinism that can lead to brain damage and death if not detected early.

Gastric dumping syndrome is encountered in approximately 25-50% of patients
25 following gastric surgery and may persist post-operatively for several months. Early dumping usually involves gastro-intestinal and vasomotor complaints. Late dumping predominantly involves vasomotor complaints and is a consequence of a reactive hypoglycaemia resulting from hyperinsulinism and an exaggerated release of glucagon-like peptide-1. The gastric dumping syndrome is infrequently

reported in children, but is difficult to diagnose and manage and has significant morbidity.

It is critical that the hypoglycaemia, secondary to the hyperinsulinism and characteristic of clinical conditions like those described above, be managed quickly if death or profound neurologic impairment is to be avoided.

The mainstay of therapy in the management of severe hypoglycaemia is glucose or dextrose infusion. In cases of drug overdose, this may follow gut decontamination. Glucagon is used sometimes but considerable caution must be taken because its success depends on limited hepatic glycogen stores. Surgical intervention is quite successful for many insulinomas where the lesion can be appropriately localised. In the case of congenital hyperinsulinism, partial or complete pancreatectomy is often necessary. Surgical excision of injection sites in cases of massive, non-accidental insulin overdose also is sometimes the preferred option.

Diazoxide is sometimes used to combat hyperinsulinism. It is a potent antihypertensive agent and acts to promote blood glucose level by suppressing insulin secretion from the pancreas. However, this drug is not specific in its treatment of hyperinsulinism and a number of undesirable side effects may follow its use. Hypotension, nausea, vomiting, dizziness, weakness and mild liver damage have been reported with its use. Severe hypoglycaemia also persists in some patients following diazoxide therapy.

Octreotide is a peptide analog of somatostatin and, like diazoxide, one of its actions is to inhibit insulin secretion from the pancreas. Octreotide has shown some promise in treating patients with hyperinsulinism, however, resistance to the drug has been reported in some patients with insulinoma. It has been used with some success in treating gastric dumping syndrome in patients refractory to standard therapy. However, its long term use is limited by side-effects such as diarrhoea and steatorrhoea. Moreover, octreotide may in some instances worsen existing hypoglycaemia by suppressing glucagon and growth hormone in the presence of unresponsive pancreatic hyperinsulinism. Octreotide therapy may also have undesired effects on reducing long term growth in infants.

Thus, as can be deduced from the foregoing discussion, the only drugs available for treating hyperinsulinism with secondary hypoglycaemia, act by suppressing further insulin secretion from the pancreas. No other drugs are presently available.

- 5 Antagonists of insulin action would be therapeutically useful agents for treating a range of diseases or clinical conditions involving hyperinsulinism and hypoglycaemia. They would work by directly competing with insulin for binding to the insulin receptor (the first step of insulin action) and would thereby counter the effects of hyperinsulinism. Their effect would be to reduce the hypoglycaemic
10 action of insulin evident in clinical conditions described above.

In view of the foregoing, it will be appreciated that there is a current on going need for biological modifiers that are capable of mimicing insulin activity.

Summary of the Invention

- The present invention relates to the use of at least a non-peptidyl compound as a
15 biological modulator of insulin activity or insulin-related activity, which compound possesses ionic and hydrophobic chemical moieties spatially located so as to mimic at least an ionic and hydrophobic amino acid residue of insulin, which amino acids are associated with the binding of insulin to its receptor.

- Compounds of the present invention exert their effects by mimicing amino acids
20 spatially located on insulin, enabling those compounds to bind to the insulin receptor or a like receptor causing biological modulation of the activity of the receptor. Compounds used in the present invention may act either as agonists or antagonists of insulin or insulin-like activity.

- Thus, in a first embodiment the invention resides in a method for treating a patient
25 suffering from one or more insulin related ailments, which method comprises the step of: administering to a patient an therapeutically effective amount of a compound that is a biological modulator of insulin activity, which compound possesses ionic and hydrophobic chemical moieties spatially located so as to mimic at least an ionic and hydrophobic amino acid residue of insulin, which
30 amino acids are associated with the binding of insulin to its receptor.

In a second embodiment, the present invention resides in the use of at least a non-peptidyl compound, which is at least a biological modulator of insulin activity and possess ionic and hydrophobic chemical moieties spatially located so as to mimic at least an ionic and hydrophobic amino acid residue of insulin, which amino acids are associated with the binding of insulin to its receptor; in the preparation of a medicament for the treatment of a patient suffering from one or more insulin related ailments.

In a third embodiment, there is provided a method for identifying a non-peptidyl compound possessing ionic and hydrophobic chemical moieties spatially located so as to mimic particular ionic and hydrophobic amino acid residues of insulin which are associated with the binding of insulin to its receptor, said method comprising the steps of: (1) comparing the three dimensional structure of the non-peptidyl compound with a three dimensional pharmacophore of an active site of insulin; and (2) selecting a non-peptidyl compound with ionic and hydrophobic chemical moieties spatially located so as to mimic said site.

In a fourth embodiment, there is provided a method for determining whether a non-peptidyl compound identified by the third embodiment of the invention is an agonist or an antagonist, said method comprising the step of: exposing the compound to an insulin or insulin like receptor and measuring the change in biological activity following exposure of the compound to the receptor.

The present invention also contemplates novel chemical compounds identified by the method of the present invention as well as the use of those molecules in a pharmaceutical composition. The pharmaceutical composition comprising a chemical compounds capable of modulating the biological activity of insulin or and a pharmaceutically acceptable carrier and/or diluent.

Brief Description of the Figures

Figure 1 shows a receptor binding pharmacophore of insulin.

Figure 2 tabulates a series of hypotheses generated from the insulin pharmacophore of Figure 1.

Figure 3 tabulates features and tolerances of the hypotheses of Figure 2 used in the searching of three dimensional chemical databases.

Figure 4 tabulates hypotheses and tolerances found to identify IM 175 in database searches, where the tolerance sphere for all features of the listed
5 hypotheses was the size indicated in the body of the table.

Figure 5 illustrates dose response of assays versus insulin concentration, showing typical dose response curves of Glucose transport (+), PTK assay (x) and Autophosphorylation experiments (o), which are expressed as a percent of maximal assay activation over a range of insulin concentrations.

10 Figure 6 summarises the effect of IM 129 on 125 I insulin bound to human placental plasma membranes (o) that is expressed as a fraction of total 125 I insulin added, and where each data point is the mean of a triplicate determination.

Figure 7 demonstrates the effect of IM 129 on 125 I insulin bound to WGA.IR (o) which is expressed as a fraction of total 125 I insulin added and overlayed with a 1-
15 site fit to the data (solid line), and where each data point is the mean of a triplicate determination.

Figure 8 illustrates the effect of IM 129 on total (in the presence of 50 nM insulin) (+) and basal (absence of insulin) (o) 32 P incorporation into FYF peptide expressed as a percentage of total 32 P- γ -ATP added to reactions, where each
20 data point is the mean of a duplicate determination \pm one standard deviation.

Figure 9 shows the effect of IM 129 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) 3 H-deoxyglucose uptake by 3T3L1 cells, where glucose transport is expressed as a percentage of a maximal 100 nM dose of insulin, and where each point is the mean of a triplicate determination \pm one standard
25 deviation.

Figure 10 represents a proposed model of antagonist competition with insulin for binding to insulin receptors, in which two alpha-beta halves of the insulin receptor joined by disulfide bonds are illustrated, activated receptors are phosphorylated

on their β -subunits. insulin is illustrated by oblong shaped objects and in which antagonist molecules are illustrated by black circular objects.

Figure 11 shows the effect of IM 129 on ^{125}I IGF-1 bound to human placental plasma membranes (o) which is expressed as a fraction of total ^{125}I IGF-1 added and is overlayed with a 1-site fit to the data (solid line) and where each data point is the mean of a triplicate determination.

Figure 12 illustrates the effect of IM 025 on ^{125}I insulin bound to human placental plasma membranes (o) which is expressed as a fraction of total ^{125}I insulin added, and where each data point is the mean of a triplicate determination.

Figure 13 summarises the effect of IM 025 on ^{125}I insulin bound to CHO.T11 cells (o) which is expressed as a fraction of total ^{125}I insulin added and overlayed with a 1-site fit to the data (solid line), and where each data point is the mean of a triplicate determination.

Figure 14 illustrates the effect of IM 025 on ^{125}I insulin bound to WGA.IR (o) which is expressed as a fraction of total ^{125}I insulin added and overlayed with a 1-site fit to the data (solid line), and where each data point is the mean of a triplicate determination.

Figure 15 demonstrates ^{32}P incorporation into FYF peptide where the effect of IM 025 on total (+) and basal (o) ^{32}P incorporation into FYF peptide is expressed as a percentage of total ^{32}P - γ -ATP added to reactions, and where each data point is the mean of a duplicate determination.

Figure 16 represents the effect of IM 025 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells expressed as a percentage of total ^3H -deoxyglucose added to cells, where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 17 shows the effect of IM 071 on ^{125}I insulin bound to human placental plasma membranes (o) expressed as a fraction of total ^{125}I insulin added where each data point is the mean of a triplicate determination.

Figure 18 summarises the effect of IM 127 on ^{125}I insulin bound to human placental plasma membranes (+) expressed as a fraction of total ^{125}I insulin added, where each data point is the mean of a triplicate determination.

Figure 19 shows the effect of IM 127 on ^{125}I insulin bound to WGA.IR (+) expressed as a fraction of total ^{125}I insulin added and overlayed with a 1-site fit to the data (solid line), where each data point is the mean of a triplicate determination.

Figure 20 illustrates ^{32}P incorporation into FYF peptide in the presence of IM 127, where the effect of IM 127 on total (in the presence of 50 nM insulin) (+) and basal (absence of insulin) (o) ^{32}P incorporation into FYF peptide is expressed as a percentage of total ^{32}P - γ -ATP added to reactions, and where each data point is the mean of a duplicate determination.

Figure 21 demonstrates the effect of IM 127 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells, where glucose transport in is expressed as a percentage of a maximal 100 nM dose of insulin and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 22 represents the effect of IM 132 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a percentage of total ^{125}I insulin added, where each data point is the mean of a triplicate determination \pm one standard deviation.

Figure 23 shows the effect of IM 132 on ^{125}I insulin bound to WGA.IR (o), expressed as a fraction of total ^{125}I insulin added and overlayed with a 1-site fit to the data (solid line), where each data point is the mean of a triplicate determination.

Figure 24 represents the effect of IM 132 on total (in the presence of 50 nM insulin) (+) and basal (absence of insulin) (o) ^{32}P incorporation into FYF peptide, expressed as a percentage of total ^{32}P - γ -ATP added to reactions, where each data point is the mean of a duplicate determination \pm one standard deviation.

- 5 Figure 25 shows the effect of IM 132 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells, where glucose transport in is expressed as a percentage of a maximal 100 nM dose of insulin, and where each point is the mean of a triplicate determination \pm one standard deviation.
- 10 Figure 26 shows the effect of IM 134 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a percentage of total ^{125}I insulin added and where each point is the mean of a triplicate determination \pm one standard deviation.

- Figure 27 demonstrates the effect of IM 134 on total (in the presence of 400 nM insulin) (o) and basal (absence of insulin) (o) ^{32}P incorporation into FYF peptide, expressed as a percentage of total ^{32}P - γ -ATP added to reactions, where two separate experiments are shown by open and solid symbols, and where each data point is the mean of a duplicate determination \pm one standard deviation.
- 15

- Figure 28 shows the effect of IM 134 on ^3H -deoxyglucose uptake in the presence (+) and absence (o) of 2 nM insulin by 3T3L1 adipocyte cells, where glucose transport in is expressed as a percentage of a maximal 100 nM dose of insulin, and each point is the mean of a triplicate determination \pm one standard deviation.
- 20

- Figure 29 illustrates the effect of IM 143 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a percentage of total ^{125}I insulin added and where each point is the mean of a triplicate determination \pm one standard deviation.
- 25

Figure 30 represents the effect of IM 143 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells, where glucose

transport is expressed as a percentage of a maximal 100 nM dose of insulin, and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 31 shows the effect of IM 144 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a percentage of total ^{125}I insulin added and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 32 illustrates the effect of IM 144 on ^{125}I insulin bound to WGA.IR (o), expressed as a fraction of total ^{125}I insulin added and overlayed with a 1-site fit to the data (solid line), in which each data point is the mean of a triplicate determination.

Figure 33 represents the effect of IM 144 on total (in the presence of 50 nM insulin) (+) and basal (absence of insulin) (o) ^{32}P incorporation into FYF peptide, expressed as a percentage of total ^{32}P - γ -ATP added to reactions, where each data point is the mean of a duplicate determination.

Figure 34 shows the effect of IM 144 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells, in which glucose transport in is expressed as a percentage of a maximal 100 nM dose of insulin, and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 35 illustrates the effect of IM 145 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a percentage of total ^{125}I insulin added and where each point is the mean of a triplicate determination.

Figure 36 shows the effect of IM 145 on total (in the presence of 50 nM insulin) (+) and basal (absence of insulin) (o) ^{32}P incorporation into FYF peptide, expressed as a percentage of total ^{32}P - γ -ATP added to reactions, in which each data point is the mean of a duplicate determination \pm one standard deviation.

Figure 37 demonstrates the effect of IM 145 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells, where glucose transport is expressed as a percentage of a maximal 100 nM dose of insulin, and each point is the mean of a triplicate determination \pm one standard deviation.

Figure 38 represents the effect of IM 171 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a percentage of total ^{125}I insulin added and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 39 shows the effect of IM 171 on total (in the presence of 400 nM insulin) (+) and basal (absence of insulin) (o) ^{32}P incorporation into FYF peptide, expressed as a percentage of total ^{32}P - γ -ATP added to reactions, where each data point is the mean of a duplicate determination \pm one standard deviation.

Figure 40 illustrates the effect of IM 171 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells, where glucose transport is expressed as a percentage of a maximal 100 nM dose of insulin, and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 41 shows the effect of IM 172 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a percentage of total ^{125}I insulin added and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 42 shows the effect of IM 172 on total (in the presence of 400 nM insulin) (+) and basal (absence of insulin) (o) ^{32}P incorporation into FYF peptide, expressed as a percentage of total ^{32}P - γ -ATP added to reactions, where each data point is the mean of a duplicate determination \pm one standard deviation.

Figure 43 illustrates the effect of IM 172 on total (in the presence of 400 nM insulin) (+) and basal (absence of insulin) (o) ^{32}P incorporation into WGA. IR 90

kDa protein band, expressed as a percentage of total ^{32}P - γ -ATP added to reactions.

Figure 44 shows the effect of IM 172 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells, in which glucose transport in is expressed as a percentage of a maximal 100 nM dose of insulin, and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 45 demonstrates the effect of IM 140 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a fraction of total ^{125}I insulin added, where each data point is the mean of a triplicate determination.

Figure 46 represents the effect of IM 140 on ^{125}I insulin bound to WGA.IR (o), expressed as a fraction of total ^{125}I insulin added, where each data point is the mean of a triplicate determination.

Figure 47 shows the effect of IM 140 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells expressed as a percentage of total ^3H -deoxyglucose added to cells, where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 48 shows the effect of IM 140 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells, where glucose transport is expressed as a percentage of a maximal 100 nM dose of insulin, and where each point is the mean of a triplicate determination \pm one standard deviation.

Figure 49 illustrates the effect of IM 140 on ^{125}I insulin binding (+) and ^3H -deoxyglucose (o) in 3T3L1 adipocyte cells, where each isotherm is expressed as a percentage of buffer control and each point is the mean of a triplicate determination \pm one standard deviation.

Figure 50 represents a possible mechanism for agonist activation of insulin receptors, in which two alpha-beta halves of the insulin receptor, joined by

disulfide bonds are illustrated, activated receptors are phosphorylated on their β -subunits, insulin is represented by oblong shaped objects and agonist molecules are represented by dumbbell shaped objects.

Figure 51 demonstrates the effect of IM 140 on ^{32}P incorporation into FYF peptide in three experiments, expressed as a percentage of maximal ^{32}P incorporation (200 nM dose of insulin) into FYF peptide.

Figure 52 demonstrates the effect of IM 140 on the total (in the presence of 100 nM insulin) (+) and basal (absence of insulin) (o) ^{32}P incorporation into FYF peptide, expressed as a percentage of total ^{32}P - γ -ATP added to reactions and where data points are expressed as the mean of a duplicate determination. Immunoprecipitated WGA.IR was used as the source of tyrosine kinase activity.

Figure 53 illustrates the effect of IM 140 on blood glucose levels in streptozotocin induced diabetic mice. The scatter plot shows the effect of IM 140 at 21 $\mu\text{mol/kg}$ (+), 30 $\mu\text{mol/kg}$ (o), 37 $\mu\text{mol/kg}$ (x) and human insulin at 2U/kg over a period of 40 minutes. A significant change in blood glucose levels in an unpaired 1-tail t-test are indicated by a single asterisk where $p < 0.05$, a double asterisk where $p < 0.01$ and a triple asterisk *** where $p < 0.001$.

Figure 54 illustrates the effect of IM 140 on ^{125}I IGF-1 bound (o) and ^{125}I insulin bound (+) to human placental plasma membranes expressed as a fraction of total ^{125}I IGF-1 and ^{125}I insulin added respectively, where each data point is the mean of a triplicate determination.

Figure 55 illustrates the effect of IM 140 on ^{125}I IGF-1 bound (x) to WGA.IR expressed as a percent of total ^{125}I IGF-1 added to assay tubes, where each data point is the mean of a triplicate determination \pm one standard deviation.

Figure 56 shows the effect of IM 175 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a fraction of total ^{125}I insulin added and overlaid with a 1-site fit to the data (solid line).

Figure 57 represents the effect of IM 175 on ^{125}I IGF-1 bound to human placental plasma membranes (o), expressed as a fraction of total ^{125}I IGF-1 added and overlaid with a 1-site fit to the data (solid line), where each data point is the mean of a triplicate determination.

- 5 Figure 58 is illustrative of the effect of IM 175 on total (presence of 2 nM insulin) (+) and basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells expressed as a percentage of total ^3H -deoxyglucose added to cells, where each data point is the mean of a triplicate determination \pm one standard deviation.

- 10 Figure 59 demonstrates the incorporation of ^{32}P into the 90 kDa protein band of immunoprecipitated insulin receptors, where the uppermost autoradiograph demonstrates phosphorylation of immunoprecipitated insulin receptor β -subunit in response to the indicated concentrations of IM 175 and insulin. Untreated lanes show phosphorylation in the absence of both insulin and IM 175, where both autoradiographs were taken from the same experiment, the leftmost (A) being a
15 48 hour exposure of the gel to X-ray film, and the rightmost (B) a 24 hour exposure. The lowermost plot demonstrates the effect of IM 175 on ^{32}P incorporation into the 90 kDa protein band of immunoprecipitated insulin receptors in the presence (o) and absence (x) of 5 nM insulin compared to a 100 nM dose of insulin (+).

- 20 Figure 60 demonstrates the effect of IM 175 on the incorporation of ^{32}P into the 90 kDa protein band of immunoprecipitated IGF-1 receptors in the presence (+) and absence (x) of 2 nM IGF-1.

- 25 Figure 61 shows the effect of IM 103 on ^{125}I insulin bound to human placental plasma membranes (o), expressed as a fraction of total ^{125}I insulin added, where each data point is the mean of a triplicate determination.

Figure 62 illustrates the effect of IM 103 on ^{125}I insulin bound to CHO.T11 cells (o) expressed as a fraction of total ^{125}I insulin added, where each data point is the mean of a triplicate determination.

Figure 63 shows the effect of IM 103 on ^{125}I insulin bound to WGA.IR (+) expressed as a fraction of total ^{125}I insulin added, where each data point is the mean of a triplicate determination \pm one standard deviation.

Figure 64 shows the effect of IM 103 on total (presence of 2 nM insulin) (+) and
5 basal (no insulin) (o) ^3H -deoxyglucose uptake by 3T3L1 cells expressed as a percentage of total ^3H -deoxyglucose added to cells, where each point is the mean of a triplicate determination.

Figure 65 illustrates The effect of IM 103 on ^{125}I IGF-1 bound to human placental
10 plasma membranes (o) expressed as a fraction of total ^{125}I IGF-1 added and overlaid with a 1-site fit to the data (solid line), where each data point is the mean of a triplicate determination.

Detailed Description of the Invention

The present invention relates to the use of at least a non-peptidyl compound as a biological modulator of insulin activity or insulin-related activity, which compound
15 possesses ionic and hydrophobic chemical moieties spatially located so as to mimic at least an ionic and hydrophobic amino acid residue of insulin, which amino acids are associated with the binding of insulin to its receptor.

In a first embodiment, the invention resides in a method for treating a patient suffering from one or more insulin related ailments, which method comprises the
20 step of: administering to a patient an therapeutically effective amount of a compound that is a biological modulator of insulin activity, which compound possesses ionic and hydrophobic chemical moieties spatially located so as to mimic at least an ionic and hydrophobic amino acid residue of insulin, which amino acids are associated with the binding of insulin to its receptor.

25 In a second embodiment, the present invention resides in the use of at least a non-peptidyl compound, which is at least a biological modulator of insulin activity and possess ionic and hydrophobic chemical moieties spatially located so as to mimic at least an ionic and hydrophobic amino acid residue of insulin, which amino acids are associated with the binding of insulin to its receptor; in the

preparation of a medicament for the treatment of a patient suffering from one or more insulin related ailments.

The phrase "biological modulator" as used herein refers to a compound that is capable of least varying insulin activity when introduced into a biological system
5 (eg an *in vitro* or *in vivo* system). Such a compound may either be an antagonist or an agonist of biological activity. Most preferably the non-peptidyl compound used in either the first or second embodiment of the invention is an insulin agonist.

- As used herein the term "patient" refers to any animal that may be suffering from one or more insulin related ailments. Most preferably the animal is a mammal.
- 10 The term "mammal" as used herein refers to a class of vertebrates whose young feed upon milk from the mother's breast. Most species are more or less hairy, all have a diaphragm, and all (except the monotremes) are viviparous. The term will be understood to include for example human, farm animals (i.e., cattle, horses, goats, sheep and pigs), household pets (i.e., cats and dogs) and the like.
- 15 The phrase "therapeutically effective amount" as used herein refers to an amount of a non-peptidyl compound sufficient to modulate a biological activity associated with the interaction of insulin with its receptors.

Insulin related ailments include ailments which are related to decreased secretion of insulin, decreased responsiveness of cells to insulin, or increased secretion of
20 insulin and may, for example, include ailments such as:- diabetes mellitus (types 1 and 2), insulinomas, insulin and hypoglycaemic drug overdose, gastric dumping syndrome and congenital hyperinsulinism. Other ailments will be known to those of ordinary skill in the field. For example, insulin and glucose therapy leads to improved cognition in alzheimers disease patients.

- 25 Amino acids associated with the binding of insulin to its receptor include, by way of example:- A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16 Tyr, A2 Ile, A3 Val and A1 Gly. As used herein the "A" or "B" nomenclature refers to either the A or B chain that forms insulin, while the number identifies the amino acid number in either chain.

While the non-peptidyl compounds employed in the present invention may mimic any number of the amino acids associated with the binding of insulin to its receptor, preferably they mimic ionic and hydrophobic moieties associated with at least two of these amino acids. More preferably, the compounds mimic ionic and hydrophobic residues on at least 4 of the abovementioned amino acids, including at least one amino acid from the group comprising:- A21 Asn, B21 Glu and A17 Glu. Even more preferably, the compounds mimic ionic and hydrophobic residues on at least 4 amino acids, including at least one amino acid selected from the group comprising A17 Glu, B21 Glu and A21 Asn, and at least one amino acid selected from the group comprising:- B24 Phe, B25 Phe, A19 Tyr, B12 Val and B12 Tyr. Desirably, the non-peptidyl compound mimics ionic and hydrophobic residues associated with at least one of the following groups of amino acid residues:-

- 1 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe;
- 15 2 A21 Asn, B21 Glu, B24 Phe, B25 Phe;
- 3 A21 Asn, B21 Glu, B24 Phe, B25 Phe, A1 Gly, A2 Ile, A3 Val;
- 4 A21 Asn, B21 Glu, A17 Glu, A19 Tyr, A1 Gly, A2 Ile, A3 Val;
- 5 A21 Asn, B21 Glu, A17 Glu, B12 Val, A1 Gly, A2 Ile, A3 Val;
- 6 A21 Asn, B21 Glu, B12 Val, A1 Gly, A2 Ile, A3 Val;
- 20 7 A21 Asn, B21 Glu, A17 Glu, B16 Tyr, A1 Gly, A2 Ile, A3 Val;
- 8 A21 Asn, B21 Glu, A17 Glu, A19 Tyr, B12 Val, B16 Tyr;
- 9 A21 Asn, B21 Glu, A19 Tyr, B12 Val, B16 Tyr;
- 10 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
- 25 11 A21 Asn, B21 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
- 12 A21 Asn, B21 Glu, B24 Phe, B25 Phe, B12 Val, B16 Tyr;
- 13 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;
- 14 A21 Asn, B21 Glu, B24 Phe, B25 Phe, A19 Tyr;
- 15 A21 Asn, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;

- 16 B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;
 17 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, B12 Val;
 18 A21 Asn, B21 Glu, B24 Phe, B25 Phe, B12 Val;
 19 A21 Asn, A17 Glu, B24 Phe, B25 Phe, B12 Val;
 5 20 B21 Glu, A17 Glu, B24 Phe, B25 Phe, B12 Val;
 21 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
 22 A21 Asn, B21 Glu, B24 Phe, B25 Phe, B16 Tyr;
 23 A21 Asn, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
 24 B21 Glu, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
 10 25 A21 Asn, B21 Glu, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
 26 A21 Asn, B21 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
 27 A21 Asn, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
 28 B21 Glu, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
 29 A21 Asn, B21 Glu, A17 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
 15 30 A21 Asn, B21 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
 31 A21 Asn, A17 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr; or
 32 B21 Glu, A17 Glu, B25 Phe, A19 Tyr, B12 Val.

Any compound capable of mimicing the spatial arrangements of the foregoing amino acids may be employed in the present invention. Preferably, the non-peptidyl compound(s) has the following formula:



where A is W or VXW;

V is V₁ or V₂;

V is substituted with up to two X groups;

V_1 is a phenyl or 6 membered heteroaromatic ring, optionally substituted with up to 5 R_1 groups, for example but not being limited to benzene, pyridine, pyridazine, pyrimidine, pyrazine, triazine;

5 V_2 is a 5 member ring system which may incorporate up to 4 hetero atoms which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, for example but not being limited to cyclopenta-1,3-diene, pyrrole, furan, thiophene, oxazole, isoxazole, pyrazole, imidazole, thiazole, isothiazole or triazole, the ring system being optionally substituted with up to 4 R_1 groups;

10 W is W_1 or W_2 or W_3 ;

W is substituted with up to two X groups;

W_1 is V_1 ;

15 W_2 is a fused bicyclic ring system comprising rings of 5 or 6 atoms, which may incorporate up to 4 hetero atoms, which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, the system being optionally substituted with up to seven R_1 groups and examples include, but are not limited to naphthalene, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, indole, benzothiophene, benzofuran, benzimidazole, indazole, 20 benzoxazole, benzisooxazole, benzthiazole, benzisothiazole, purine, indoline, isoindoline;

W_3 is $-N(R_2)R'_2$;

R_1 is independently H, OH, alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy, haloalkyl, haloalkoxy, halogen, SH, thioalkyl, cyano (-CN), 25 $N(R_2)R'_2$, phenyl, phenyl optionally substituted with up to five alkyl groups of 1 to 3 carbon atoms or up to five halogen atoms, benzyl, phenethyl, nitro, -COR₃, -R₅COR₃, -R₅SOR₃, -R₅SO₂R₃, -SO₂N(R₂)R'₂ or azido;

30 R_2 and R'_2 are independently H, alkyl of 1 to 6 carbon atoms, alkenyl of 3 to 6 carbon atoms, alkynyl of 3 to 6 carbons, hydroxyalkyl of 2 to 6 carbons, alkoxy of 2 to 6 carbons, haloalkyl, haloalkenyl, haloalkoxy, benzyl, benzyl

optionally substituted with up to four R_1 groups, phenylethyl, phenylethyl optionally substituted with up to four R_1 groups, arylalkyl, and where R_2 and R'_2 can also be joined to form cyclic structures including, but not limited to pyrrolidine, piperidine, hexahydro-1H-azepine, morpholine or piperazine;

5 R_3 is independently H, OH, alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy, $-R_4N(R_2)R'_2$, mesyl, triflouromesyl, $-NHSO_2CH_3$ or $-NHSO_2CF_3$;

R_4 is independently a bond, alkyl, alkenyl or alkynyl;

10 X is independently, a bond, $-R_4N(R_2)R_4-$, $-R_4N=NR_4-$, $-R_4N(R_2)-N(R_2)R_4-$, $-R_4OR_4-$, $-R_4SR_4-$, $-R_5-$, $-R_5O-$, $-R_5S-$, $-R_5N(R_2)-$, $-SO-$, sulfonyl ($-SO_2-$), $-CO-$, $-CONH-$, $-NHCONH-$, $-NHCO-$, $-CONHCO-$, $-CON(R_2)-$, $-R_5COR_5-$, $-R_5COR_5N(R_2)R_5-$, $-N(R_2)CO-$ or $-R_4N(R_2)R_4COR_4-$;

R_5 is independently alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy;

Y is either Y_1 , Y_2 or Y_3 ;


15 Y is substituted with at least two, but optionally up to four X linking groups;
 Y_1 is a fused bicyclic ring system comprising rings of 5 or 6 atoms which may incorporate up to 4 hetero atoms, which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, the ring system optionally independently incorporating a sulfoxide (SO), sulfone (SO₂) or carbonyl (CO) group and optionally up to seven R_1 groups, for
20 example but not limited to croman, isochroman, benzofuran, cromene, 1,2,3,4-tetrahydronaphthalene, 1,4-dihydronaphthalene, indan, indene, benzopiperidine, indoline, isoindoline, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline or pteridine, coumarin or
25 2,3-dihydrocoumarin;

Y_2 is a 6:6:6 or a 6:5:6 fused tricyclic system which may incorporate up to 4 hetero atoms which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, the ring system optionally independently incorporating a sulfoxide (SO), sulfone (SO₂) or carbonyl (CO)
30 group, and the ring system being substituted with at least two, but optionally

up to four X linking groups and optionally up to seven R_1 groups and thus examples include, but are not limited to 9H-xanthone, 9H-xanthene, phenoxathiin, phenoxathiin-10-oxide, phenoxathiin-10-dioxide, acridine, phenazine, phenothiazine, phenoxazine, phenothiazine-5-oxide, phenothiazine-5-dioxide, thiathrene-5-dioxide, thiathrene-5-oxide, carbazole, dibenzo[b,d]furan, dibenzo[b,d]thiophene;

Y_3 is V_1 ;

Z is independently $-R_6COOH$, $-R_6SO_3H$, $-R_6NO_2$, $-R_6SO_2H$, $-R_6SO_2NHR_2$; $-R_7SO_2NHCOR_4$, $-N$ -trifluoromesylsulfonamidate, $-OH$, -2 -yl-hydroxyethanoic acid ($-CH(OH)COOH$), -3 -yl-2-hydroxypropanoic acid ($-CH_2CH(OH)COOH$) - 2 -yl-2-hydroxypropanoic acid ($-CH(CH_3)(OH)COOH$), -3 -yl-2,3-dihydroxypropanoic acid ($-CH(OH)CH(OH)COOH$), -2 -yl-2,3-dihydroxypropanoic acid ($-C(CH_2(OH))(OH)COOH$), -3 -yl-2-hydroxypropan-3-one-1-oic acid ($-COCH(OH)COOH$), 2 -yl-2-hydroxypropandioic acid ($-C(COOH)(OH)COOH$), -2 -yl-propandioic acid ($-C(COOH)(H)COOH$), -4 -yl-2-hydroxybutan-4-one-1-oic acid ($-COCH_2CH(OH)COOH$), 2 -yl-2-hydroxybutan-1,4-dioic acid ($-C(OH)(COOH)CH_2COOH$), 3 -yl-2-hydroxybutan-1,4-dioic acid ($-CH(CH(OH)COOH)COOH$), 5 -yl-tetrazole,



R_6 is independently a bond, alkyl, alkenyl, alkynyl, alkoxy, $-CO(CH_2)_n-$, where n is an integer between 0 and 4, alkanolic, alkenolic or alkynolic;

with the exception that, where W_1 is an optionally substituted phenyl then Y_1 cannot be an optionally substituted phenyl.

The term "alkyl" as used herein refers to an alkane derived radical of between 1 and 6 carbon atoms unless otherwise defined, including straight, branched or cyclic alkane derived radicals, monovalent or bivalent alkane derived radicals in that it may be joined to one or two groups via any allowed bond to its carbon atoms. Preferably, the straight or branched alkyl groups are methyl, ethyl, propyl,

isopropyl, butyl, *i*-butyl, *t*-butyl, pentyl or hexyl and the cycloalkyl groups are cyclopentyl or cyclohexyl.

The term "alkenyl" as used herein refers to a hydrocarbon derived radical containing from 2 to 6 carbon atoms, unless otherwise defined and at least one carbon to carbon double bond, and includes straight, branched or cyclic hydrocarbon derived radicals, monovalent or bivalent hydrocarbon derived radicals in that such alkane derived radicals may be joined to one or two groups via any allowed bond to its carbon atoms. Preferably, the alkenyl groups are ethenyl, propenyl, butenyl, cyclopentenyl, or cyclobutenyl.

- 10 The term "alkynyl" as used herein refers to a hydrocarbon derived radical containing from 2 to 6 carbon atoms and at least one carbon to carbon triple bond and includes straight chained or branched hydrocarbon derived radicals, monovalent or bivalent hydrocarbon derived radicals in that such hydrocarbon derived radicals may be joined to one or two groups via any allowed bond to its
- 15 carbon atoms. Preferably, the alkynyl groups are ethynyl, propynyl or butynyl.

The term "alkoxy" as used herein refers to an alkyl group of indicated carbon atoms, attached to other groups through an oxygen linkage (monovalent) or; to other groups through an oxygen linkage and a bond to any of its allowed carbon atoms (bivalent).

- 20 The term "hydroxyalkoxy" as used herein refers to an alkoxy group substituted at one or more carbon atoms, with one or more hydroxyl groups.

The term "haloalkoxy" as used herein refers to an alkoxy group substituted at one or more carbon atoms, with one or more halogen atoms. Preferably, the halogen substituents are fluorine, chlorine, bromine or iodine.

- 25 The term "alkanol" as used herein refers to an alkyl group substituted at one or more carbon atoms, with one or more hydroxyl groups.

The term "haloalkyl" as used herein refers to an alkyl group substituted at one or more carbon atoms, with one or more halogen atoms. Preferably, the halogen substituents are fluorine, chlorine, bromine or iodine.

The term "thioalkyl" as used herein refers to an alkyl group of indicated carbon atoms, attached to other groups through a sulfur linkage.

The term "alkanoic" as used herein refers to an alkyl group of indicated carbon atoms, substituted at one carbon atom with a carboxylic acid group (-COOH).

- 5 The term "alkenoic" as used herein refers to an alkenyl group of indicated carbon atoms, substituted at one carbon atom with a carboxylic acid group (-COOH).

The term "alkynoic" as used herein refers to an alkynyl group of indicated carbon atoms, substituted at one carbon atom with a carboxylic acid group (-COOH).

- 10 More preferably, when the non-peptidyl compound employed in the invention has the structure of formula 1 above and V is V_1 or V_2 , then:

V_1 is selected from the group consisting of, benzene, pyridine, pyridazine, pyrimidine, pyrazine or triazine and is optionally substituted with up to 5 R_1 groups; and

- 15 V_2 is selected from the group consisting of, cyclopenta-1,3-diene, pyrrole, furan, thiophene, oxazole, isoxazole, pyrazole, imidazole, thiazole, isothiazole or triazole and is optionally substituted with up to 4 R_1 groups;

and W is W_2 then

- 20 W_2 is selected from the group consisting of naphthalene, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, indole, benzothiophene, benzofuran, benzimidazole, indazole, benzoxazole, benzisooxazole, benzthiazole, benzisothiazole, purine, indoline or isoindoline and is optionally substituted with up to seven R_1 groups;

and Y is either Y_1 or Y_2 then

- 25 Y_1 is selected from the group consisting of croman, isochroman, benzofuran, cromene, 1,2,3,4-tetrahydronaphthalene, 1,4-dihydronaphthalene, indan, indene, benzopiperidine, indoline, isoindoline, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline or pteridine, coumarin or 2,3-dihydrocoumarin and
30 is optionally substituted with up to seven R_1 groups; and

Y₂ is selected from the group consisting of 9H-xanthone, 9H-xanthene, phenoxathiin, phenoxathiin-10-oxide, phenoxathiin-10-dioxide, acridine, phenazine, phenothiazine, phenoxazine, phenothiazine-5-oxide, phenothiazine-5-dioxide, thiathrene-5-dioxide, thiathrene-5-oxide, 5 carbazole, dibenzo[b,d]furan or dibenzo[b,d]thiophene and is optionally substituted with up to seven R₁ groups.

Desirably, when the non-peptidyl compound employed in the invention has the structure of formula 1 above and A is W or VXW then:

V is phenyl or pyrazole, optionally substituted with up to 5 R₁ groups;
10 and W is W₁, W₂ or W₃ then
W₁ is phenyl optionally substituted with up to 5 R₁ groups;
W₂ is naphthalene or quinoline optionally substituted with up to seven R₁ groups wherein R₁ is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;
15 W₃ is -N(R₂)R₂ wherein R₂ is propyl;
X is independently, a bond, methoxy (-OCH₂-), oxypropoxy (-O(CH₂)₃O-), hexenyloxy (-O(CH₂)₄CH=CH-), sulfonyloxy (-SO₂O-), methyl (-CH₂-), amidyl (-CONH-) or -NHCONH-;
and Y is either Y₁ or Y₂ then
20 Y₁ is croman, 4-H-chromen-4-one or naphthalene optionally substituted with up to seven R₁ groups wherein R₁ is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;
Y₂ is 9H-xanthone optionally substituted with up to seven R₁ groups
25 wherein R₁ is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;
Y₃ is phenyl optionally substituted with up to 5 R₁ groups wherein R₁ is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl; and
30 Z is independently -R₆COOH, -R₆SO₃H or -N-trifluoromesylsulfonamidate wherein R₆ is independently a bond or propyl.

By way of example, the non-peptidyl compound(s) are selected from the group:

1. 4,4'-Methylenebis[3-hydroxy-2-naphthalenecarboxylic acid]
(herein referred to as IM 025);
2. 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-
5 1-benzopyran-2-carboxylic acid
(herein referred to as IM 071);
3. 2,4-dichloro-6-(*N*-(trifluoromethanesulfonyl))sulfamoylphenyl 3,5-dichloro-2-
hydroxybenzenesulfonate
(herein referred to as IM 103);
- 10 4. 7-[(4-acetyl-3-hydroxy-2-propylphenyl)methoxy]-4-oxo-8-propyl-4*H*-1-
benzopyran-2-carboxylic acid.
(herein referred to as IM 127);
5. 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-
2*H*-1-benzopyran-2-carboxylic acid
15 (herein referred to as IM 129);
6. 3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-(1*H*-pyrazol-3-
yl)phenoxy]propyl]oxy]-2*H*-1-benzopyran-2-carboxylic acid
(herein referred to as IM 132);
7. 3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-ethoxyphenoxy]propyl]oxy]-
20 2*H*-1-benzopyran-2-carboxylic acid
(herein referred to as IM 134);
8. 3-[4-[7-carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-
hydroxyphenoxy]propoxy]-9*H*-xanthene]]propanoic acid
(herein referred to as IM 140);
- 25 9. 8-propyl-7-(quinol-2'-ylmethoxy)-3,4-dihydro-2*H*-1-benzopyran-2-carboxylic
acid
(herein referred to as IM 143);

10. 7-(naphth-2'-ylmethoxy)-8-propyl-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid

(herein referred to as IM 144);

11. N-(trifluoromethanesulfonyl)-3,5-dinitro-4-(N,N'-dipropylamino)benzenesulfonamide

(herein referred to as IM 145);

12. 8-propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid

(herein referred to as IM 171);

13. 3,4-dihydro-7-[[6-(4-methoxyphenyl)hexenyl]oxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid

(herein referred to as IM 172); or

14. 8,8'-[Carbonylbis[imino-3,1-phenylenecarbonylimino(4-methyl-3,1-phenylene)carbonylimino]]bis-1,3,5-naphthalenetrisulfonic acid

(herein referred to as IM 175).

The non-peptidyl compounds described herein may also be prepared as dimers or heterodimers of compounds of the above mentioned formula $AXYZ_n$, where such compounds are joined through a X linking group by way of their V or W groups.

It will be understood by those of ordinary skill in the field that the compounds of the present invention may also be prepared as dimers and/or heterodimers. Furthermore analogues and precursors of the abovementioned compounds may also be prepared for use in the invention.

The compounds of the present invention include chemical derivatives that may be converted to the above mentioned compounds *in vivo*, such derivatives including but not being limited to esters and amides.

The present invention also contemplates pharmaceutical composition comprising at least a chemical compound capable of modulating the biological activity of insulin and a pharmaceutically acceptable carrier and/or diluent.

The compounds selected for use in the invention are desirably prepared in a purified form suitable for administration to a patient. Purification of such compounds may be achieved by any means known in the art, such as distillation, chromatographic means *etc.*

- 5 Once purified the compounds may be formulated into therapeutics as neutral or salt forms. Pharmaceutically acceptable salts include, for example, the acid addition salts (formed with any free amino groups of the compounds) and which are formed with inorganic acid such as hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, maleic and the like. Salts formed with free
- 10 acidic groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or iron (III) hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like. The composition may further comprise excipients that are pharmaceutically acceptable and compatible with the active ingredient. Examples
- 15 of excipients which may be used in such a formulation include water, saline, ethanol, dextrose, glycerol or the like, or combinations thereof. If desired, the composition may also contain minor amounts of auxiliary substances such as wetting agents, pH buffering agents, or emulsifying agents which enhance the compounds.
- 20 Pharmaceutical compositions may be administered by injection, or prepared for oral, pulmonary, nasal or for any other form of administration. Preferably the pharmaceutically acceptable composition(s) are administered, for example, intravenously, subcutaneously, intramuscularly, intraorbitally, ophthalmically, intraventricularly, intracranially, intracapsularly, intraspinally, intracisternally,
- 25 intraperitoneally, buccal, rectally, vaginally, intranasally or by aerosol administration.

The mode of administration must, however, be at least suitable for the form in which the composition has been prepared. The mode of administration for the most effective response may need to be determined empirically and the means of

30 administration described below are given as examples, and do not limit the method of delivery of the composition of the present invention in any way. All the

above formulations are commonly used in the pharmaceutical industry and are commonly known to suitably qualified practitioners.

The compounds provided herein can be formulated into pharmaceutical compositions by admixture with pharmaceutically acceptable nontoxic excipients and carriers and administered by any parenteral techniques such as
5 subcutaneous, intravenous and intraperitoneal injections. In addition the formulations may optionally contain one or more adjuvants.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions (where water-soluble) or dispersions and sterile powders for the
10 extemporaneous preparation of sterile injectable solutions or dispersion. Alternatively, the compounds of the invention may be encapsulated in liposomes and delivered in injectable solutions to assist their transport across cell membrane. Alternatively or in addition such preparations may contain constituents of self-assembling pore structures to facilitate transport across the cellular membrane.
15 The carrier may be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use
20 of surfactants. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions may be prepared by incorporating the active compounds in the required amount in an appropriate solvent with various of the
25 other ingredients enumerated above, as required, followed by filtered sterilisation. Generally, dispersions are prepared by incorporating the various sterilised active ingredient into a sterile vehicle that contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of
30 preparation are vacuum drying and freeze-drying techniques that yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

Oral Delivery

Contemplated for use herein are oral solid dosage forms, which are described generally in *Martin, Remington's Pharmaceutical Sciences*, 18th Ed. (1990 Mack Publishing Co. Easton PA 18042) at Chapter 89, which is herein incorporated by
5 reference. Solid dosage forms include tablets, capsules, pills, troches or lozenges, cachets or pellets. Also, liposomal or proteinoid encapsulation may be used to formulate the present compositions (as, for example, proteinoid microspheres reported in U.S. Patent No. 4,925,673). Liposomal encapsulation may be used and the liposomes may be derivatised with various polymers (*E.g.*, U.S. Patent No.
10 5,013,556). A description of possible solid dosage forms for the therapeutic is given by Marshall, in *Modern Pharmaceutics*, Chapter 10, Banker and Rhodes ed., (1979), herein incorporated by reference. In general, the formulation will include the compounds described as part of the invention (or a chemically modified form thereof), and inert ingredients which allow for protection against the stomach
15 environment, and release of the biologically active material in the intestine.

For the compounds described as part of the invention the location of release may be the stomach, the small intestine (the duodenum, the jejunum, or the ileum), or the large intestine. One skilled in the art has available formulations that will not dissolve in the stomach, yet will release the material in the duodenum or elsewhere in the
20 intestine. Preferably, the release will avoid the deleterious effects of the stomach environment, either by protection of the composition or by release of the compounds beyond the stomach environment, such as in the intestine.

To ensure full gastric resistance, a coating impermeable to at least pH 5.0 is essential. Examples of the more common inert ingredients that are used as enteric
25 coatings are cellulose acetate trimellitate (CAT), hydroxypropylmethylcellulose phthalate (HPMCP), HPMCP 50, HPMCP 55, polyvinyl acetate phthalate (PVAP), Eudragit L30D, Aquateric, cellulose acetate phthalate (CAP), Eudragit L, Eudragit S, and Shellac. These coatings may be used as mixed films.

A coating or mixture of coatings can also be used on tablets, which are not intended
30 for protection against the stomach. This can include sugar coatings, or coatings that make the tablet easier to swallow. Capsules may consist of a hard shell (such as

gelatin) for delivery of dry therapeutic *i.e.* powder; for liquid forms, a soft gelatin shell may be used. The shell material of cachets could be thick starch or other edible paper. For pills, lozenges, molded tablets or tablet triturates, moist massing techniques can be used.

- 5 The therapeutic can be included in the formulation as fine multiparticulates in the form of granules or pellets of particle size about 1mm. The formulation of the material for capsule administration could also be as a powder, lightly compressed plugs or even as tablets. The therapeutic could be prepared by compression.

- 10 Colourants and flavoring agents may all be included. For example, compounds may be formulated (such as by liposome or microsphere encapsulation) and then further contained within an edible product, such as a refrigerated beverage containing colorants and flavoring agents.

- One may dilute or increase the volume of the therapeutic with an inert material. These diluents could include carbohydrates, especially mannitol, alpha-lactose, anhydrous lactose, cellulose, sucrose, modified dextrans and starch. Certain
15 inorganic salts may be also be used as fillers including calcium triphosphate, magnesium carbonate and sodium chloride. Some commercially available diluents are Fast-Flo, Emdex, STA-Rx 1500, Emcompress and Avicell.

- Disintegrants may be included in the formulation of the therapeutic into a solid
20 dosage form. Materials used as disintegrants include but are not limited to starch including the commercial disintegrant based on starch, Explotab. Sodium starch glycolate, Amberlite, sodium carboxymethylcellulose, ultramylopectin, sodium alginate, gelatin, orange peel, acid carboxymethyl cellulose, natural sponge and bentonite may all be used. Another form of the disintegrants are the insoluble
25 cationic exchange resins. Powdered gums may be used as disintegrants and as binders and these can include powdered gums such as agar, Karaya or tragacanth. Alginic acid and its sodium salt are also useful as disintegrants.

- Binders may be used to hold the therapeutic compounds together to form a hard tablet and include materials from natural products such as acacia, tragacanth, starch
30 and gelatin. Others include methylcellulose (MC), ethyl cellulose (EC) and

carboxymethyl cellulose (CMC). Polyvinyl pyrrolidone (PVP) and hydroxypropylmethyl cellulose (HPMC) could both be used in alcoholic solutions to granulate the therapeutic.

5 An antifrictional agent may be included in the formulation of the therapeutic to prevent sticking during the formulation process. Lubricants may be used as a layer between the therapeutic and the die wall, and these can include but are not limited to: stearic acid including its magnesium and calcium salts, polytetrafluoroethylene (PTFE), liquid paraffin, vegetable oils and waxes. Soluble lubricants may also be used such as sodium lauryl sulfate, magnesium lauryl sulfate, polyethylene glycol of
10 various molecular weights, and Carbowax 4000 and 6000.

Glidants that might improve the flow properties of the compound during formulation and to aid rearrangement during compression might be added. The glidants may include starch, talc, pyrogenic silica and hydrated silicoaluminate.

To aid dissolution of the therapeutic into the aqueous environment, a surfactant
15 might be added as a wetting agent. Surfactants may include anionic detergents such as sodium lauryl sulfate, dioctyl sodium sulfosuccinate and dioctyl sodium sulfonate. Cationic detergents might be used and could include benzalkonium chloride or benzethonium chloride. The list of potential nonionic detergents that could be included in the formulation as surfactants are lauromacrogol 400, polyoxyl
20 40 stearate, polyoxyethylene hydrogenated castor oil 10, 50 and 60, glycerol monostearate, polysorbate 40, 60, 65 and 80, sucrose fatty acid ester, methyl cellulose and carboxymethyl cellulose. These surfactants could be present in the formulation of the compounds either alone or as a mixture in different ratios.

Additives which potentially enhance uptake of the compounds are for instance the
25 fatty acids oleic acid, linoleic acid and linolenic acid.

Controlled release formulation may be desirable. The compounds could be incorporated into an inert matrix that permits release by either diffusion or leaching mechanisms *i.e.*, gums. Slowly degenerating matrices may also be incorporated into the formulation. Another form of a controlled release of this therapeutic is by a
30 method based on the Oros therapeutic system (Alza Corp.), *i.e.* the drug is enclosed

in a semipermeable membrane which allows water to enter and push drug out through a single small opening due to osmotic effects. Some enteric coatings also have a delayed release effect.

- A mix of materials might be used to provide the optimum film coating. Film coating
5 may be carried out in a pan coater or in a fluidized bed or by compression coating.

Pulmonary Delivery

Also contemplated herein is pulmonary delivery of the compounds. The compounds may be delivered to the lungs of a mammal while inhaling and traverses across the lung epithelial lining to the blood stream.

- 10 Contemplated for use in the practice of this invention are a wide range of mechanical devices designed for pulmonary delivery of therapeutic products, including but not limited to nebulizers, metered-dose inhalers, and powder inhalers, all of which are familiar to those skilled in the art.

- Some specific examples of commercially available devices suitable for the practice
15 of this invention are the Ultravent nebulizer, manufactured by Mallinckrodt, Inc., St. Louis, Missouri; the Acorn II nebulizer, manufactured by Marquest Medical Products, Englewood, Colorado; the Ventolin metered dose inhaler, manufactured by Glaxo Inc., Research Triangle Park, North Carolina; and the Spinhaler powder inhaler, manufactured by Fisons Corp., Bedford, Massachusetts.

- 20 All such devices require the use of formulations suitable for the dispensing of the compounds. Typically, each formulation is specific to the type of device employed and may involve the use of an appropriate propellant material, in addition to the usual diluents, adjuvants and/or carriers useful in therapy. Also, the use of liposomes, microcapsules or microspheres, inclusion complexes, or other types of
25 carriers is contemplated.

Formulations suitable for use with a nebulizer, either jet or ultrasonic, will typically comprise the compounds suspended in water. The formulation may also include a buffer and a simple sugar (e.g., for protein stabilization and regulation of osmotic pressure). The nebulizer formulation may also contain a surfactant, to reduce or

prevent surface induced aggregation of the compounds caused by atomization of the solution in forming the aerosol.

Formulations for use with a metered-dose inhaler device will generally comprise a finely divided powder containing the compounds suspended in a propellant with the aid of a surfactant. The propellant may be any conventional material employed for this purpose, such as a chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethanol, and 1,1,1,2-tetrafluoroethane, or combinations thereof. Suitable surfactants include sorbitan trioleate and soya lecithin. Oleic acid may also be useful as a surfactant.

Formulations for dispensing from a powder inhaler device will comprise a finely divided dry powder containing the compound and may also include a bulking agent, such as lactose, sorbitol, sucrose, or mannitol in amounts which facilitate dispersal of the powder from the device, e.g., 50 to 90% by weight of the formulation. The compounds (or derivative) should most advantageously be prepared in particulate form with an average particle size of less than 10 microns, most preferably 0.5 to 5 microns, for most effective delivery to the distal lung.

Nasal delivery of the compounds is also contemplated. Nasal delivery allows the passage of the protein to the blood stream directly after administering the therapeutic product to the nose, without the necessity for deposition of the product in the lung. Formulations for nasal delivery include those with dextran or cyclodextran.

Compounds of the present invention should be administered in dosage unit form that is prophylactically and/or therapeutically effective. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the novel dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active material for the treatment sought. Thus the quantity of active compound to be

administered will be largely dependent on, the toxicity and specific activity of compound, the subject to be treated and the degree of treatment required. Precise amounts of compound required to be administered may depend on the judgement of the practitioner and may be peculiar to each subject.

- 5 It will be appreciated that the composition may be given as a single dose schedule, or preferably, in a multiple dose schedule. A multiple dose schedule is one in which a primary course of delivery may be with 1 to 10 separate doses, followed by other doses given at subsequent time intervals required to maintain or reinforce the treatment. The dosage regimen will also, at least in part, be
10 determined by the need of the individual and the judgement of the practitioner.

In addition, the composition of the present invention containing the non-peptidyl compounds may be administered in conjunction with other agents and compounds, for example, immunoglobulins, sulfonylureas, biguanides, α -glucosidase inhibitors, thiazolidinediones, diazoxide, octreotide, insulin
15 secretagogues as appropriate.

According to a third embodiment, there is provided a method for identifying a non-peptidyl compound possessing ionic and hydrophobic chemical moieties spatially located so as to mimic particular ionic and hydrophobic amino acid residues of insulin which are associated with the binding of insulin to its receptor, said method
20 comprising the steps of: (1) comparing the three dimensional structure of the non-peptidyl compound with a three dimensional pharmacophore of an active site of insulin; and (2) selecting a non-peptidyl compound with ionic and hydrophobic chemical moieties spatially located so as to mimic said site.

Preferably, the active site comprises at least four amino acid residues selected
25 from the group A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16 Tyr, A2 Ile, A3 Val and A1 Gly, with at least one amino acid residue being selected from the group A21 Asn, B21 Glu and A17 Glu. Desirably, the active site comprises one of the following groups of amino acid residues:-

- 1 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe;
- 30 2 A21 Asn, B21 Glu, B24 Phe, B25 Phe;

- 3 A21 Asn, B21 Glu, B24 Phe, B25 Phe, A1 Gly, A2 Ile, A3 Val;
4 A21 Asn, B21 Glu, A17 Glu, A19 Tyr, A1 Gly, A2 Ile, A3 Val;
5 A21 Asn, B21 Glu, A17 Glu, B12 Val, A1 Gly, A2 Ile, A3 Val;
6 A21 Asn, B21 Glu, B12 Val, A1 Gly, A2 Ile, A3 Val;
5 7 A21 Asn, B21 Glu, A17 Glu, B16 Tyr, A1 Gly, A2 Ile, A3 Val;
8 A21 Asn, B21 Glu, A17 Glu, A19 Tyr, B12 Val, B16 Tyr;
9 A21 Asn, B21 Glu, A19 Tyr, B12 Val, B16 Tyr;
10 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16
Tyr;
10 11 A21 Asn, B21 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
12 A21 Asn, B21 Glu, B24 Phe, B25 Phe, B12 Val, B16 Tyr;
13 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;
14 A21 Asn, B21 Glu, B24 Phe, B25 Phe, A19 Tyr;
15 A21 Asn, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;
15 16 B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;
17 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, B12 Val;
18 A21 Asn, B21 Glu, B24 Phe, B25 Phe, B12 Val;
19 A21 Asn, A17 Glu, B24 Phe, B25 Phe, B12 Val;
20 B21 Glu, A17 Glu, B24 Phe, B25 Phe, B12 Val;
20 21 A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
22 A21 Asn, B21 Glu, B24 Phe, B25 Phe, B16 Tyr;
23 A21 Asn, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
24 B21 Glu, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
25 A21 Asn, B21 Glu, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
25 26 A21 Asn, B21 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
27 A21 Asn, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
28 B21 Glu, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
29 A21 Asn, B21 Glu, A17 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
30 A21 Asn, B21 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;

31 A21 Asn, A17 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr; or

32 B21 Glu, A17 Glu, B25 Phe, A19 Tyr, B12 Val.

Comparison of the three dimensional structure of the non-peptidyl compound with the three dimensional pharmacophore may involve comparison of a minimum
5 energy structure of the non-peptidyl compound with a three dimensional pharmacophore of the active site of insulin, the position of the amino acid residues comprising the active site being determined with reference to crystal and/or NMR solution structures of insulin and its analogues. The three dimensional pharmacophore may incorporate tolerance spheres about the positions of the
10 respective amino acids to reflect the conformational flexibility of insulin.

The positions of the amino acid residues of the active site are defined relative to one another by the following three dimensional coordinates (in hundredths of Angstroms) within tolerance spheres of between 1.5 and 5 Å: B12 Val (343.9, 157.2, -242.6); B25 Phe (113, -5.3, 780.8); B24 Phe (92.9, 569.7, 43.2); B16 Tyr
15 (361.8, 1068, -467.5); B21 Glu (-84.9, 1711.2, 99.6); A17 Glu (-1079.2, 1005.6, -67.9); A19 Tyr (-763.8, 49.2, -4.1); A21 Asn (-344.2, 908, 677.9); A1 Gly (-870.2, -574.9, 127.9); A2 Ile (-489.8, -30, -218.1); and A3 Val (-237.6, -703.3, -348.4).

In a more specific embodiment of the present invention, the tolerance spheres for each residue are as follows: 5Å for residues B24 Phe, B25 Phe, A17 Glu, B21
20 Glu, A21 Glu, A1 Gly, A2 Ile; 3.6 Å for residues A19 Tyr, B16 Tyr, B12 Val and 3Å for A3 Val.

An efficient means to select a non-peptidyl insulin mimetic compound from a potentially large number of non-peptidyl compounds involves comparing non-peptidyl compounds against a three dimensional pharmacophore of insulin using
25 a computer program, for example *Catalyst* (MSI), to screen one or more computerised databases of three dimensional chemical structures of non-peptidyl compounds.

If, for example, the *Catalyst* program is used, the amino acids in the active site are preferably represented by the following *Catalyst* defined features: B12 Val -
30 hydrophobic; B25 Phe - hydrophobic; B24 Phe - hydrophobic; B16 Tyr -

hydrophobic; B21 Glu - negative ionisable; A17 Glu - negative ionisable; A19 Tyr - hydrophobic; A21 Asn - negative ionisable; A1 Gly - positive ionisable; A2 Ile - hydrophobic; and A3 Val - hydrophobic.

Preferably still, the amino acid residues represented within *Catalyst* as
5 hydrophobic may be further specified as: B12 Val - aliphatic; B25 Phe - aromatic; B24 Phe - aromatic; B16 Tyr - aromatic; A19 Tyr - aromatic; A2 Ile - aliphatic; and A3 Val - aliphatic.

According to a fourth embodiment, there is provided a method for determining whether a non-peptidyl compound identified by the third embodiment of the
10 invention is an agonist or an antagonist, said method comprising the step of: exposing the compound to an insulin or insulin like receptor and measuring the change in biological activity following exposure of the compound to the receptor.

Best Method of Performing the Invention

Further features of the present invention will be more fully described in the
15 following Examples. It is to be understood, however, that this detailed description is included solely for the purposes of exemplifying the invention, and should not be understood in any way as a restriction on the broad description as set out above.

Pharmacophore Construction and Use for Database Searching

20 Models of Insulin used for pharmacophore construction

Many crystal and NMR-solution structures of insulin and its analogues have been published. The NMR solution structure pdb1hiu.ent was chosen from the Brookhaven protein database to allow for flexibility of the insulin molecule in solution. Additionally, because the insulin crystal structure is reported to be an
25 inactive "closed" conformation and that an "open" conformation, represented by glycine[B24] insulin, is active, the NMR solution structure, pdb1hit, of Gly[B24]insulin was also chosen for pharmacophore construction. The pdb1hiu.ent file contained a collection of eleven models (MODEL 1 through MODEL 11) while the glycine mutant contained nine models (MODEL 0 through

MODEL 8). Each structure represented a possible solution conformation of the molecule that satisfies the NMR distance data.

Model Analysis

The eleven native insulin models (HIU 1-11) were overlaid with each other and an average structure was determined. This structure was processed by fixing the alpha carbons and subjecting it to 100 steps of Steepest Descent minimisation (using CHARM force field) to normalise the slightly distorted coordinates of the rest of the molecule. This same procedure could not be used with the glycine analog due to the larger variation among the structures; therefore each model was extracted as a separate molecular structure file. Since the pharmacophore is based upon the native structure of insulin, each glycine model was re-mutated back to Phenylalanine at position B24 using QUANTA's Protein Design facility. The conformation of the phenylalanine side chain was adjusted to minimise bad contacts and the side chain rotamer most likely to be found by Karplus rules was assigned. The structures were then analysed for their degree of "openness" by measuring the distance between a pseudo atom (placed between residues A2 and A3) and residue B25. The average native insulin structure and two glycine mutant structures HIT3 and HIT4 were chosen to represent a range of "openness" in insulin conformations.

In QUANTA, the following residues from each model structure were selected: A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16 Tyr, A2 Ile, A3 Val and A1 Gly. The remainder of the molecule was removed and residues were connected with virtual "bonds". The coordinates of these pseudo structures were then imported into the *Catalyst* (MSI) software program and pharmacophore construction was conducted by mapping features (negative ionisable, positive ionisable, hydrophobic) onto the residues. Original structures were then overlayed onto the hypothesis to estimate tolerance spheres about each feature. The pharmacophore is shown in Figure 1. The pharmacophore was then used to create subsets of features (Hypotheses) that were used to screen databases. Figure 2 shows the thirteen working hypotheses generated - including variations used for database searching. A maximum 5Å tolerance was assigned where the variation of side chains about a feature was large. The

tolerances about each feature are shown in Figure 3. Smaller 3Å tolerance spheres were used for all HIT3 and HIT4 models. In subsequent searches all features were given identical tolerance spheres and the size of these were adjusted from a maximum of 5Å to a minimum of 1.5Å to limit the number and quality of compounds identified.

These hypotheses were then used to rapidly screen chemical databases - that contained representative three-dimensional structures of each compounds likely minimum energy conformations, for those compounds that contained specific functional groups that could be positioned in the tolerance spheres of each hypothesis.

Results

Hypotheses 2 and 7a, with the tolerances about each of the features of the hypotheses (negative ionisable, positive ionisable and hydrophobic) as listed in Figure 3 identified the following compounds:-

| Compound | Ref. used herein |
|--|------------------|
| 4,4'-Methylenebis[3-hydroxy-2-naphthalenecarboxylic acid] | IM 025 |
| 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid | IM 071 |
| 2,4-dichloro-6-(N-(trifluoromethanesulfonyl)sulfamoylphenyl 3,5-dichloro-2-hydroxybenzenesulfonate | IM 103 |
| 7-[(4-acetyl-3-hydroxy-2-propylphenyl)methoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid. | IM 127 |
| 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid | IM 129 |
| 3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-(1H-pyrazol-3-yl)phenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid | IM 132 |
| 3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-ethoxyphenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid | IM 134 |
| 3-[4-[7-carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]-9H-xanthene]]propanoic acid | IM 140 |
| 8-propyl-7-(quinol-2'-ylmethoxy)-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid | IM 143 |
| 7-(naphth-2'-ylmethoxy)-8-propyl-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid | IM 144 |

| | |
|--|--------|
| N-(trifluoromethanesulfonyl)-3,5-dinitro-4-(<i>N,N</i> -dipropylamino)benzenesulfonamide | IM 145 |
| 8-propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid | IM 171 |
| 3,4-dihydro-7-[[6-(4-methoxyphenyl)hexenyl]oxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid | IM 172 |
| 8,8'-[Carbonylbis(imino-3,1-phenylenecarbonylimino(4-methyl-3,1-phenylene)carbonylimino)]bis-1,3,5-naphthalenetrisulfonic acid | IM 175 |

IM 175 was identified by a number of hypotheses. The number of hypotheses that identified IM 175 depended upon the tolerances of features within each hypothesis. Figure 4 shows the multiple hypotheses that identified compound IM 175 and the tolerances of features in each hypothesis.

- 5 For each of the identified compounds, tests were undertaken to demonstrate their ability to compete for insulin binding in a variety of insulin receptor preparations.

Further tests were undertaken to ascertain their ability to stimulate the insulin receptor second messenger system and their ability to effect insulin's stimulation of its second messenger system. This was determined in three assays: the
 10 compounds ability to stimulate: insulin receptor autophosphorylation; phosphorylation of a synthetic peptide that is a substrate of the insulin receptor β -subunit associated tyrosine kinase activity and glucose transport in 3T3L1 adipocyte cells. Furthermore, *in vivo* tests were undertaken to ascertain their ability to lower blood glucose levels in experimentally induced diabetic mice.

15 Solubility of Compounds

Compounds were solubilised in a minimal volume of DMSO before diluting into the relevant assay buffer. Final assay concentrations of DMSO were kept to less than 1% (volume/volume) and assays were controlled for any effect of the residual DMSO.

- 20 For *in vivo* studies compounds were dissolved in a sterile solution of 1.6% glycerin, 0.25% m-cresol, pH 6.75.

Insulin Binding Assay

The Hepes buffer system used to dilute the compound depended on the source of insulin receptor preparation used for each assay:

Human Placental Plasma Membranes - 50 mM Hepes, pH 7.8 @ 4 °C containing
5 100 U/mL Bacitracin; CHO cells transfected with human insulin receptor (CHO.T11) - Hams media supplemented with 25 mM Hepes, pH 7.8@ 4°C and 0.1% BSA; Wheat germ agglutinin affinity purified insulin receptors, insulin affigel affinity purified receptors - 50 mM Hepes, pH 7.8 @ 4°C containing 100 U/mL Bacitracin and 0.02% Triton-X 100.

10 Where possible, compounds were prepared in the absence of DMSO because DMSO concentrations greater than 0.05% DMSO effects insulin binding to its receptors.

One hundred microlitres of compound was mixed on ice with 50 µL of assay buffer containing 5 fmole of ¹²⁵I insulin (7 x 10⁷ MBq.mmol⁻¹) and 50 µL of insulin
15 receptor solution. Tubes were mixed and incubated for 16 h at 4°C. After the incubation, the tubes were placed on ice and 300 µL of a 1:5 suspension of 0.4% bovine-γ-globulin in 20% polyethyleneglycol 6000 (BDH chemicals) was added to each tube, mixed thoroughly and left to incubate for 30 min on ice. The tubes were then centrifuged (3,000g) for 30 min at 4°C, the supernatant was aspirated
20 off and the pellet counted for radioactivity. The ability of each compound to compete for ¹²⁵I insulin binding to each of the receptor sources was compared to unlabelled human insulin. Inhibition constants were estimated by graphical analysis of displacement plots or calculated using the curve fitting programs EBDA and/or LIGAND.

25 Activation of Insulin Receptor Second Messenger System

The activity of each of the compounds was evaluated in three assays of insulin action: 1) Insulin receptor autophosphorylation; 2) Phosphorylation of a synthetic substrate specific for insulin receptor (PTK assay) and 3) glucose uptake by 3T3L1 adipocyte cells.

Insulin Receptor Phosphorylation

Insulin receptor phosphorylation was assessed using a modification of the method described by Li *et al.* (21). Insulin receptors prepared from CHO cells stably expressing full-length insulin receptors (CHO.T11) (15 μ L) were preincubated at room temperature for 20 minutes with 7.5 μ L of 50 mM HEPES, pH 7.4 containing 150 mM NaCl, 0.1 mM phenylmethylsulfonylfluoride, 10 mM MgCl₂, 2 mM MnCl₂ and 0.1% Triton X-100 and varying concentrations of human insulin or putative insulin mimetic compound. After this preincubation, 2.5 μ L of 1 mM ATP, 100 mM sodium orthovanadate and 5 μ Ci of [γ -³²P]ATP were added and incubated at room temperature for 25 minutes. The reaction was terminated by the addition of 1.5 μ L of 0.5 M EDTA and 3.5 μ L of 1.5 M NaCl. The solution was then either solubilised and resolved by SDS-PAGE or immunoprecipitated using anti-insulin R β (29B4) mouse monoclonal IgG or anti-phosphotyrosine (PY20) mouse monoclonal IgG (Santa Cruz Biotechnology Inc., USA), solubilised and then resolved by SDS-PAGE. Immunoprecipitation was performed by incubating one microgram of antibody with the reaction mixture over night at 4°C. Following this incubation, 40 μ L of protein A Sepharose CL-4B (Pharmacia Biotech AB, Sweden) in a 75%:25%(v/v) slurry with 50 mM Tris, pH 7.5 containing 150 mM NaCl (TBS) was added to the receptor-antibody mix and incubated for a further 90 minutes at 4°C with continuous stirring. The receptor-antibody-Sepharose complex was separated from the supernatant by centrifugation at 7,500 g for 5 minutes. The pellet was washed three times in TBS containing 0.1% Triton X-100 and 10 μ L of 0.1 M glycine, pH 3.3 was added. The solution was then solubilised by incubation for 5 minutes at 100°C in 10 μ L of 80 mM Tris-HCl, pH 6.8 containing 0.45 M DTT, 56 mM EDTA, 30% sucrose and 6% SDS. Following centrifugation for 5 minutes at 7,500 g, the supernatant was resolved on a 7.5% SDS polyacrylamide gel. The gel was dried under vacuum and exposed to X-ray film. In some instances, gel bands corresponding to 90 kDa were excised and the associated radioactivity was measured by suspending the gel band in 2.5 mL of Scintillation fluid (Beckman) using a Beckman β -counter.

Protein Tyrosine Kinase (PTK) Assay

Protein tyrosine kinase activity was determined using a modification of the method of N. Konstantopoulos (22). The assay involved measuring the effect of compounds on the insulin receptor mediated incorporation of ^{32}P into a specific
5 synthetic peptide substrate (RRDIFETDYFRK), (FYF peptide).

The assay was commenced by adding 5 μL of an insulin receptor solution in 50 mM Hepes, pH 7.5 to: 5 μL of the test compound in 50 mM Hepes, pH 7.5; 5 μL of FYF peptide in 40 mM imidazole pH 7.3 containing 40 mM β -glycerophosphate, 1 mM EGTA, 100 mM MgCl_2 , 5 mM MnCl_2 and 0.05% BSA; 5 μL of
10 phosphorylation mixture containing 0.1 mM ATP, 0.5 mM sodium vanadate and 0.5 μCi of $[\gamma\text{-}^{32}\text{P}]\text{-ATP}$ and; 5 μL of a 50 mM Hepes, pH 7.5 solution alone or containing 2 μM insulin. The reaction was allowed to proceed for 20 minutes at 30°C before it was terminated by the addition of 12.5 μL of 7.5 M guanidine hydrochloride. An aliquot (10 μL) of the reaction mixture was then spotted onto a
15 Whatman 81 phosphocellulose paper (15 mm x 15 mm), washed three times (2 min per wash) in about 50 mL of 30% acetic acid, 0.05% orthophosphoric acid then finally, with one wash in about 200 mL of water. The papers were then rinsed briefly (15 sec) in 95% ethanol, dried, suspended in 2.5 mL of scintillation fluid (Beckman) and counted for radioactivity in a Beckman β -counter.

20 Glucose Transport in 3T3-L1 Cells

Glucose transport was determined using a modification of the method of S.J. Isakoff *et al.* (23). The assay involved measuring the effect of compounds on the uptake of ^3H -deoxyglucose into 3T3L1 adipocyte cells.

Growth and Differentiation of 3T3-L1 Cells

25 The cell line, 3T3-L1, derived from mouse embryo fibroblasts, was grown in 250 mL tissue culture flasks containing 15 mL medium made up of DMEM (Dulbecco's modified Eagle's medium containing glutamine, 2 mM; sodium bicarbonate, 30 mM; penicillin 71.5 mg/litre; streptomycin, 152.5 mg/litre) supplemented with 10% foetal calf serum (FCS), pH 7.4. Cells were maintained in a humidified incubator

at 37°C in a 5% CO₂ environment. The medium was changed every 3 days and the cells were split when they were about 70-80% confluent.

3T3-L1 fibroblast cells from one tissue culture flask were suspended in 50 mL of DMEM media containing 10% FCS and 0.5 mL of this suspension were seeded
5 into each well of two 48-well tissue culture plates. The differentiation of the cells was initiated when the fibroblast cells were fully confluent. Routinely, differentiation was induced on cells that had been growing in tissue culture plates for 4-5 days. Cells were exposed to 0.5 mL of DMEM medium containing 10% FCS, 2 µg/mL insulin, 0.1 µg/mL dexamethasone, 0.5 mM isobutylmethylxanthine
10 and 1 µg/mL biotin for three days. Cells were then incubated a further three days in 1 mL of post differentiation DMEM medium containing 10% FCS and 2 µg/mL insulin. The cells were subsequently maintained with DMEM medium containing 10% FCS. The differentiated cells were examined microscopically between 11 and 13 days after differentiation to ensure they displayed the adipocyte
15 phenotype.

Glucose Transport Assay

One day prior to the glucose transport assay, cell medium was replaced with DMEM containing 10% charcoal-treated foetal calf serum (CTFCS). On the day of the assay, the medium was aspirated and each well was washed with 1 mL of
20 25 mM Hepes, pH 7.4 containing 0.9% NaCl (NaCl-HEPES). One hundred microlitres of NaCl-HEPES buffer in the absence and presence of 2 nM insulin and/or test compound was then added to wells and left to incubate for 20 min at 37°C. Ten microlitres of NaCl-HEPES containing 3 mM 2-deoxy-D-glucose and 0.1 µCi 2-deoxy-D-[1-³H]glucose were then added. Following a 10 min incubation
25 at 37°C, the medium was aspirated off and the wells rapidly washed three times with 1 mL of ice-cold 50 mM phosphate, pH 7.5 containing 150 mM NaCl (PBS) and 10 mM glucose. Two hundred microlitres of 3% Triton X-100 was then added to each well and left to incubate for a minimum of 10 minutes. One hundred microlitres of the solubilised cells was then added to 2.5 mL of liquid scintillation
30 fluid and counted for radioactivity in a Beckman β-counter.

Figure 5 illustrates that dose response curves of insulin are clearly different in each of the three assays used to classify compounds. These differences have been reported previously in assays that measure the signalling and metabolic actions of insulin. For example, M.A. Soos *et al.* (24) showed that the half maximal insulin concentration for stimulation of ^3H -deoxyglucose was 0.1 nM in NIH 3T3 HIR3.5 cells which is 30 fold lower than that reported for receptor phosphorylation. They also studied a number of insulin mimetic monoclonal anti-insulin receptor antibodies and showed differential stimulation of 2-deoxyglucose uptake in NIH 3T3 HIR3.5 cells (100%), [^3H]thymidine incorporation (30%) and ^{32}P incorporation into insulin receptor β -subunit (0%). However, a later publication showed that in using a more sensitive assay for receptor autophosphorylation, the same antibodies stimulate phosphorylation of the insulin receptor β -subunit (25). The differences between assays is not clearly understood but may be due in part to differences in temperature and incubation times of the experiments (24). However, it is also recognised that there are substantial differences between whole cell and broken cell preparations (25). For these reasons it is necessary to characterise each compound in a number of assays.

***In vivo* Studies**

Induction of Experimental Diabetes

Six week old male CD-1 mice were injected *intra peritoneally* with a sterile solution of streptozotocin prepared in 100 mM citrate buffer, pH 4.65 at a dose of 200 mg/kg and an injection volume of 16.7 mL/kg. Mice were housed in a 12 h light/dark cycle and fed a normal chow diet *ad libitum*. Drinking water was supplemented with 5% sucrose for the first two days following streptozotocin injection before being replaced with tap water thereafter. After 4 days, blood was taken from the saphenous vein of the mice using the method of Hem, A *et al.* (26). Random non-fasting blood glucose levels were determined on venous blood using an esprit glucometer (Bayer). Mice with blood glucose levels of greater than 15 mmol/L were characterised as diabetic.

Blood Glucose Level Measurement

Blood glucose levels of mice were determined, followed by an *intra peritoneal* dose of IM compound dissolved in diluent (1.6% glycerin, 0.25% m-cresol, pH 6.75). Mice were then bled periodically and whole blood glucose levels were
5 determine using a one-touch esprit glucometer (Bayer). The ability of compounds to lower blood glucose levels was compared at each time point to mice injected with diluent alone (negative control) and those injected with a 2U/kg *intra peritoneal* dose of human insulin (positive control).

Antagonist Compounds

10 The following compounds have been classified as antagonists of insulin action: IM 025, IM 071, IM 127, IM 129, IM 132, IM 134, IM 143, IM 144, IM 145, IM 171 and IM 172). Some of these compounds show synergism with insulin, which is detailed in individual cases below.

IM 129 shows experimental data characteristic of the other IM compounds listed
15 above. These characteristics are detailed below. All other IM antagonist compounds are included thereafter with minimum discussion, unless explanation of an extraordinary result is required.

EXAMPLE 1

7-[3-(4-Acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-
20 benzopyran-2-carboxylic acid sodium salt hydrate (IM 129)

This compound may be synthesised by the method described by S. W. Djuric *et al.* (27), incorporated herein by reference.

Effect on Insulin Binding

IM 129 competes with insulin for binding to a number of receptor sources.
25 Displacement plots from competition studies conducted using membrane-bound receptor preparations display a characteristic skewed bell-shaped isotherm with a peak in ¹²⁵I insulin binding at about 100 to 200 µM. At higher concentrations, IM 129 inhibits insulin binding with an apparent K_i between 320 µM and 550 µM. A

typical displacement plot from four separate experiments can be seen in Figure 6. In contrast, purified receptor preparations yield a less complicated displacement plot with an estimated K_i of 189 μM (*LIGAND*, 1-site fit), as can be seen in Figure 7, which is consistent with observed differences in insulin binding to membrane-associated and soluble receptor preparations described in the background art.

Effect on Biological Activity

The dose response curves of IM 129 in several biological assays indicate that it is an antagonist of insulin action with an apparent IC_{50} between 30 and 200 μM , as illustrated by Figures 8 and 9.

- 10 The difference between the apparent K_i estimated from insulin binding and biological assays appears to be an artefact of the skewed bell-shaped displacement plots obtained in binding experiments. Mechanistically, this is likely to be caused by a complex competition between the compound and insulin for the multiple binding sites on the insulin receptor. Figure 10 shows a possible model
- 15 for receptor activation that is consistent with P. De Meyts (14) and L. Schaffer (15) models, where insulin receptors are activated through cross-linking of two sites on each α -subunit as discussed earlier. Receptor deactivation occurs when cross-linking is disturbed by the antagonist, the inhibition constant of which is measured in biological assays. However, the inhibition constant measured in biological
- 20 assays correlates with an increase in bound tracer insulin in binding experiments (Bell-shaped curve), which indicates that an additional insulin molecule is accommodated in the binding site. This may occur because the receptor binding site is no longer obstructed by a cross-linked insulin molecule or a compound induced conformational change occurs. In fact, insulin itself induces just such a
- 25 change at its binding site, accommodating one molecule at physiological concentrations, a second at greater than 100 nM and even a third molecule at supra physiological concentrations of insulin. As a result of the additional tracer insulin bound, higher concentrations of compound are then required to compete off the bound insulin, leading to a higher apparent inhibition constant observed in
- 30 insulin binding assays compared with biological assays.

Specificity

IM 129 displaces ^{125}I IGF-1 binding from human placental plasma membranes (HPPM) in a dose dependent manner with an apparent K_i of 0.8-1.3 mM, as illustrated by Figure 11. This is approximately two fold higher than the apparent
5 K_i for ^{125}I insulin binding for the same source of receptors. Therefore, IM 129 has different specificity for the two receptors and is a less potent inhibitor of ^{125}I IGF-1 binding than it is for ^{125}I insulin binding.

EXAMPLE 2

4,4'-Methylenebis[3-hydroxy-2-naphthalenecarboxylic acid] (IM 025)

10 4,4'-Methylenebis[3-hydroxy-2-naphthalenecarboxylic acid] is commercially available from the Sigma Chemical Company.

Effect on Insulin Binding

IM 025 competes with insulin binding to insulin receptors from a number of sources with an estimated K_i of 100 - 400 μM , as seen in Figures 12 to 14. Figure
15 12 shows a typical competition experiment of IM 025 on ^{125}I insulin binding to human placental plasma membranes. The calculated K_i over 5 experiments is $335 \mu\text{M} \pm 34 \mu\text{M}$. Figure 13 shows a typical competition experiment of IM 025 on ^{125}I insulin binding to CHO.T11 cells. The estimated K_i is 137-337 μM (*LIGAND*, 1-site fit) over two experiments. Figure 14 shows a typical competition experiment
20 of IM 025 on ^{125}I insulin binding to WGA.IR. The estimated K_i is $108 \pm 32 \mu\text{M}$ (*LIGAND*, 1-site fit).

Effect on Biological Activity

As can be seen in Figures 16 and 17, IM 025 is an antagonist of insulin action. IM 025 causes a dose dependent decrease in the incorporation of ^{32}P into FYF
25 peptide in control (no insulin) and insulin stimulated tubes, as illustrated by Figure 15. Figure 16 shows that IM 025 also inhibits glucose transport in 3T3L1 cells.

The estimated IC₅₀ of IM 025 in these two assays is 150 μ M and 170 μ M respectively.

EXAMPLE 3

7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-
5 benzopyran-2-carboxylic acid sodium salt (IM 071)

Effect on Insulin Binding

Figure 17 shows that IM 071 competes for insulin binding to HPPM in a dose dependent manner, with an inhibition constant (calculated for two experiments) of 186 μ M \pm 33 μ M. IM 071 is structurally related to IM 129 and is likely to have
10 similar antagonistic effects of insulin in biological assays.

EXAMPLE 4

7-[(4-Acetyl-3-hydroxy-2-propylphenyl)methoxy]-4-oxo-8-propyl-4H-1-benzopyran-
2-carboxylic acid (IM 127)

7-[(4-Acetyl-3-hydroxy-2-propylphenyl)methoxy]-4-oxo-8-propyl-4H-1-benzopyran-
15 2-carboxylic acid can be obtained by the method described in European Patent Application EP 132, 366 incorporated herein by reference.

Effect on Insulin Binding

IM 127 displays a skewed bell-shaped displacement plot of ¹²⁵I insulin binding to HPPM with a peak in binding between 10 and 100 μ M and competes for insulin
20 binding with an apparent K_i of 140-320 μ M, as can be seen in Figure 18. Figure 19 shows that IM 127 competes for ¹²⁵I insulin binding to WGA.IR in a less complex manner with an apparent K_i of 295 μ M (LIGAND, 1-site fit).

Effect on Biological Activity

IM 127 displays insulin antagonistic activity in a number of assays, summarised in
25 Figures 20 and 21. Figure 21 specifically shows that the estimated K_i is 60 μ M in the glucose transport assay.

EXAMPLE 5

3,4-Dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-(1H-pyrazol-3-yl)phenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid (IM 132)

3,4-Dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-(1H-pyrazol-3-yl)phenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid can be synthesised by the method described by R. W. Harper *et al.* (28), incorporated herein by reference.

Effect on Insulin Binding

Figures 22 and 23 illustrate that IM 132 effects insulin binding with an apparent inhibition constant of 200 to 450 μ M, which is dependent upon receptor source.

Effect on Biological Activity

IM 132 is an antagonist of insulin action because it does not increase 32 P incorporation into FYF peptide or activate glucose transport. Figures 24 and 25 illustrate that it also inhibits the insulin stimulation of these assays. The estimated inhibition constant of the compound in the PTK and glucose transport assays is 360 μ M and 64 μ M respectively. However, in the presence of insulin, IM 132 shows significant synergism with insulin in the glucose transport assay, increasing the effect of a submaximal dose of insulin (2 nM) to levels above those attained by 100 nM insulin. This indicates that IM 132 may be interacting with the insulin receptor at the insulin binding site in a more complex manner than IM 129.

EXAMPLE 6

3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-ethoxyphenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid (IM 134)

3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-ethoxyphenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid can be obtained by the method described in International Application WO 95 17, 183, incorporated herein by reference.

Effect on Insulin Binding

Figure 26 shows that IM 134 competes with 125 I insulin binding to HPPM with an apparent K_i of 900 μ M.

Effect on Biological Activity

- 5 IM 134 is an antagonist of insulin action because it does not increase 32 P incorporation into FYF peptide or glucose transport and inhibits the insulin stimulation of these assays, as can be seen in Figures 27 and 28. The estimated inhibition constants of the compound in these assays are 300-400 μ M and 60-110 μ M respectively.

10

EXAMPLE 7

8-propyl-7-(quinol-2'-ylmethoxy)-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid
(IM 143)

Effect on Insulin Binding

- As Figure 29 shows, IM 143 competes with 125 I insulin binding to HPPM with an
15 apparent K_i of about 2 mM.

Effect on Biological Activity

IM 143 is an antagonist of insulin action because it does not stimulate glucose transport activity and inhibits a stimulating (2 nM) dose of insulin with an estimated inhibition constant of 40 μ M as illustrated in Figure 30.

EXAMPLE 8

7-(naphth-2'-ylmethoxy)-8-propyl-3,4-dihydro-2*H*-1-benzopyran-2-carboxylic acid (IM 144)

Effect on Insulin Binding

- 5 IM 144 effects insulin binding in a range of receptor sources, evident from Figures 31 and 32. IM 144 causes a skewed bell-shaped displacement plot of ¹²⁵I insulin binding to HPPM with an apparent *K_i* of 900 μM.

Effect on Biological Activity

- 10 IM 144 is an antagonist of insulin action because it does not increase ³²P incorporation into FYF peptide or activate glucose transport and inhibits the insulin stimulation of these assays, as shown in Figures 33 and 34. The estimated inhibition constant of the compound in the PTK and glucose transport assays is 130 μM and 125 μM respectively.

EXAMPLE 9

- 15 *N*-(trifluoromethanesulfonyl)-3,5-dinitro-4-(*N,N'*-dipropylamino)benzenesulfonamide (IM 145)

Effect on Insulin Binding

IM 145 competes for ¹²⁵I insulin binding to HPPM with an apparent *K_i* of about 1000 μM as shown in Figure 35.

- 20 **Effect on Biological Activity**

Figures 36 and 37 show that IM 145 is an antagonist of insulin action because it does not stimulate PTK or glucose transport activity and inhibits a stimulating dose of insulin (400 nM and 2 nM respectively) in each assay with an estimated IC₅₀ of 90 and 60 μM respectively.

EXAMPLE 10

8-Propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid (IM 171)

8-Propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-
5 2H-1-benzopyran-2-carboxylic acid can be obtained by the method described in International Application WO 95/17183 (Eli Lilly and Co) incorporated herein by reference.

Effect on Insulin Binding

Figure 38 shows that IM 171 competes for ¹²⁵I insulin binding to HPPM with an
10 apparent K_i of 370 μM.

Effect on Biological Activity

IM 171 is an antagonist of insulin action. It does not increase ³²P incorporation into FYF peptide or activate glucose transport and inhibits insulin stimulation of each of these assays, as is evident from Figures 39 and 40. The estimated IC₅₀
15 in the PTK and glucose transport assays are 250 μM and 80 μM respectively.

EXAMPLE 11

3,4-dihydro-7-[[6-(4-methoxyphenyl)hexenyl]oxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid (IM 172)

Effect on Insulin Binding

20 IM 172 competes for ¹²⁵I insulin binding to HPPM with an apparent K_i of 218 μM, as Figure 41 illustrates.

Effect on Biological Activity

IM 172 is an antagonist of insulin action. It does not increase ³²P incorporation into FYF peptide or the insulin receptor β-subunit and does not activate glucose
25 transport in 3T3L1 cells. As Figures 42 to 44 demonstrate, it also inhibits the

insulin stimulation of each of these assays. The estimated inhibition constant of the compound is 80 μ M in the both the PTK and glucose transport assays.

Agonist Compounds

IM 140, IM 175, IM 103 have been classified as agonists of insulin action.

5

EXAMPLE 12

3-[4-[7-carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]-9H-xanthene]]propanoic acid disodium trihydrate (IM 140)

3-[4-[7-Carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]-9H-xanthene]]propanoic acid disodium trihydrate can
10 be synthesised by the method described by J. S. Sawyer *et al.* (29), incorporated herein by reference.

Effect on Insulin Binding

The effect of IM 140 on 125 I insulin binding to insulin receptors prepared from a
15 number of sources is complex, as is evident from Figures 45 and 46. At low micromolar concentrations, IM 140 causes an increase in 125 I insulin binding with an apparent EC50 of 10-20 μ M. Maximal 125 I insulin binding occurs at concentrations of about 80-100 μ M. Figure 45 clearly shows that concentrations above 100 μ M cause a dose dependent inhibition of 125 I insulin binding with an
20 apparent Ki of between 250 and 570 μ M.

The complicated nature of the 125 I insulin binding isotherm can be explained by a similar mechanism as described for IM 129. However, IM 140 is likely to cross-link the α 1 and α 2 sites of adjacent α -subunits, because it activates the insulin receptor. In fact, as described below, IM 140 is also synergistic with insulin,
25 indicating that both insulin and IM 140 interact with the receptor in a similar manner. However, the initial increase in 125 I insulin that corresponds to the activation of receptors, indicates that IM 140 must also cause a change in the

receptor that allows additional molecules of tracer insulin to bind. This change may be analogous in mechanism to that described earlier for IM 129 and insulin.

Effect on Biological Activity

IM 140 is an agonist of insulin action because, as Figures 47 to 49 illustrate, it
5 activates glucose transport in 3T3L1 cells, as Figures 51 and 52 illustrate, it causes phosphorylation of an exogenous insulin receptor tyrosine kinase substrate by insulin receptors of differing levels of purity and, as Figure 53 demonstrates, lowers blood glucose levels in experimentally induced diabetic mice. IM 140 has an efficacy of 25-40% compared with insulin (inter experimental
10 variation) and exhibits a bi-phasic dose-response curve with an apparent EC₅₀ of 20-30 μ M (Glucose transport). Moreover, as Figure 48 and 49 demonstrate IM 140 acts in a synergistic manner with a sub-maximal dose of insulin (2 nM) to stimulate glucose transport to 80% of maximal efficacy (100 nM insulin) at a concentration of 15 μ M with an apparent EC₅₀ of 8 μ M. The observed difference
15 in concentration of IM 140 effecting insulin binding and biological activity appears to be, at least in part, an artefact of inter-assay variation. When these parameters were measured using identical receptor sources (3T3L1 adipocyte cells), identical buffers and the same temperature, IM 140 was found to effect each assay over a similar concentration range, as illustrated in Figure 49.

20 Figures 51 shows that IM 140 also promotes the ATP-dependent phosphorylation of endogenous FYF peptide in a dose dependent manner. Figure 52 indicates that when immunoprecipitated WGA.IR receptors are used as the source of tyrosine kinase activity, IM 140 promotes phosphorylation of endogenous FYF peptide in a bi-phasic manner. The immunoprecipitated WGA.IR receptors were
25 only stimulated by a 100 fold excess of IGF-1 relative to insulin, indicating that the preparation was IGF-1 receptor free. This indicates that the phosphorylation of exogenous FYF peptide substrate was via the insulin receptor.

Figure 53 demonstrates that IM140 significantly lowers blood glucose levels in diabetic mice with an efficacy of about 25% of human insulin at a dose of 35
30 μ mol/kg. Preliminary studies also indicate that IM 140 does not decrease blood glucose levels in diabetic mice at a concentration of 100 μ mol/kg. Therefore IM

140 displays a bi-phasic dose response curve *in vivo* with a peak activity of 20-40 $\mu\text{mol/kg}$ and an EC_{50} of 15 $\mu\text{mol/kg}$.

The bi-phasic dose response curve is analogous to growth hormone activation of its receptor. This indicates that IM 140 may be activating the insulin receptor by cross-linking its α -subunits. This is consistent with the model of insulin's interaction with its receptor proposed by P. De Meyts (14) and L. Schaffer (15). They propose that insulin activates the insulin receptor by cross-linking the α_1 and α_2 sites on adjacent insulin receptor α -subunits. IM 140 presumably contains some features that also enable it to cross-link these sites. The bell shaped dose response curve can then be explained by high concentrations of IM 140 binding each of the α -subunits thereby preventing cross-linking of the receptor (Figure 50). IM compounds that are antagonists would lack some of the features of IM 140 or due to size or conformational restraints may be unable to interact concomitantly with both the α_1 and α_2 sites. Antagonist compounds may be similar to growth hormone mutants that are modified at one of their receptor contact sites which leads to them becoming potent antagonists of hGH.

Specificity

IM 140 effects ^{125}I insulin and ^{125}I IGF-1 binding to HPPM in a dose dependent and complex manner, as can be seen in Figure 54. However, over the range of IM 140 tested, the compound does not compete for ^{125}I IGF-1 binding to HPPM. In contrast, at high concentrations IM 140 promotes ^{125}I IGF-1 binding to HPPM which may be a shift to the right of the bell shaped displacement curve observed for ^{125}I insulin binding. Figure 55 indicates that IM140 does not compete for ^{125}I IGF-1 binding in WGA purified receptor preparations from solubilised CHO wild-type cells. These cells are almost devoid of insulin receptors but have high endogenous populations of IGF-1 receptors.

It is not surprising that IM 140 effects both IGF-1 binding and Insulin binding to membrane preparations considering the homology and degree of cross-reactivity between the hormones and their receptors. Each receptor probably has a common shaped binding domain that can accommodate either hormone in a

structurally equivalent manner. This result suggests that IM 140 has features that allow it to interact more specifically with the insulin receptor than the IGF-1 receptor, within a common binding domain.

EXAMPLE 13

- 5 8,8'-[Carbonylbis[imino-3,1-phenylenecarbonylimino(4-methyl-3,1-phenylene)carbonylimino]]bis-1,3,5-naphthalenetrisulfonic acid hexasodium salt (IM 175)

Effect on Insulin Binding and Specificity

- IM 175 competes with ^{125}I insulin for binding to HPPM. Figure 56 shows that the
10 apparent K_i is 46-48 μM . IM 175 competes for binding according to a simple 1-site model and does not display the complex binding kinetics associated with other agonist compounds (eg IM 140). This may be because IM 175 is a larger compound and its size may prevent additional molecules of insulin interacting with the receptor. IM 175 also competes for ^{125}I IGF-1 binding to HPPM with an
15 estimated K_i of 9 μM as shown in Figure 57), suggesting that it is a more potent competitor of IGF-1 binding than insulin binding. These results support the premise that both insulin and IGF-1 receptors have a common hormone binding site and also suggests that IM compounds are accommodated within this site. Differential specificity of compounds for either receptor is explained by structural
20 differences between compounds.

Effect on Biological Activity

IM 175 promotes glucose transport to levels approaching maximal stimulation by insulin in 3T3L1 adipocyte cells. Figure 58 shows a typical experiment, the apparent EC_{50} is about 250 μM .

- 25 Figure 59 demonstrates that IM 175 stimulates autophosphorylation of insulin receptor β -subunits with about 65% of the efficacy of a 100 nM dose of insulin. The immunoprecipitated insulin receptors used for this experiment are free of contaminating IGF-1 receptors, indicating that IM 175 specifically stimulates the phosphorylation of the insulin receptor. Figure 60 illustrates that IM 175 can also

stimulate the IGF-1 receptor, where immunoprecipitated IGF-1 receptors are free of insulin receptors. However, the dose-response curves indicate that IM175 has a lower potency on IGF-1 receptors than it does on insulin receptors.

IM 175 is structurally a symmetrical molecule and is a dimer of other IM compounds. It fits a general formula of ZXYXA-AXYXZ whereas IM 140 fits the general formula AXYXZ. This dimeric structure may be related to it having the highest efficacy of agonist compounds as other studies have implicated a role of symmetry and dimer structure in increased molecule activity. For example, small peptide agonists of the erythropoietin receptor and thrombopoietin receptor, when dimerised, activate their respective receptors with greater potency than monomers. Additionally, bivalent anti insulin receptor antibodies activate insulin receptors, whereas monovalent Fab fragments do not.

EXAMPLE 14

2,4-dichloro-6-(*N*-(trifluoromethanesulfonyl)sulfamoylphenyl 3,5-dichloro-2-hydroxybenzene sulfonate (IM 103)

Effect on Insulin Binding

IM 103 competes with insulin for binding to insulin receptors in a dose dependent manner. The displacement plot is similar to other IM compounds (exemplified by IM 129 and IM 140) and does not fit a simple 1-site or 2-site model. The apparent inhibition constant is dependent upon receptor source and is $137 \pm 45 \mu\text{M}$ (*LIGAND*, 1-site fit) in HPPM, $132 \pm 29 \mu\text{M}$ in CHO.T11 cells and $83 \mu\text{M}$ in WGA.IR receptors, as can be seen in Figures 61, 62 and 63 respectively.

Effect on Biological Activity

IM 103 is an agonist of insulin action displaying a bi-phasic biological dose response curve with an apex at concentration of $110 \mu\text{M}$ and an apparent EC_{50} of $45 \pm 7 \mu\text{M}$, as illustrated by Figure 64. The efficacy is about 15% of that achieved at a maximal dose of insulin (100 nM). IM 103 abolishes the insulin stimulation of glucose transport in a dose dependent manner. However the dose response curve indicates complex kinetics.

Specificity

IM 103 displaces ^{125}I IGF-1 binding from HPPM in a dose dependent manner with an apparent K_i of 59 μM (*LIGAND* estimated 1-site fit), as can be seen in Figure 65 which compares with a K_i for ^{125}I insulin binding of $137 \pm 45 \mu\text{M}$ (*LIGAND* estimated 1-site fit). Thus, IM 103 appears to have a higher specificity for IGF-1 receptors than insulin receptors.

Modifications and variations such as would be apparent to the skilled addressee are considered to fall within the scope of the present invention.

References

1. Hubbard *et al.* Nature (1994) 372 746-54;
2. Hubbard Embo J. (1997) 16 5572-81
3. Garrett *et al.* Nature (1998) 394 395-99
4. Lou *et al.* Science (1999) 285 1077-80
5. Kahn *et al.* J. Biol. Chem. (1974) 249 2249-2257
6. Hammond *et al.* Am. J. Physiol. (1997) 272 E1136-44
7. De Meyts *et al.* Biochem. Biophys. Res. Commun. (1973) 55 154-161
8. Helmerhorst Biochem. Biophys. Res. Commun. (1987) 147 399-407
9. Pollet *et al.* J. Biol. Chem. (1977) 252 5828-5834
10. Donner Proc. Natl. Acad. Sci. USA (1980) 77 3176-3180
11. Corin and Donner J. Biol. Chem. (1982) 257 104-110
12. Helmerhorst and Yip Biochemistry (1993) 32 2356-62
13. Donner and Yonkers J. Biol. Chem. (1983) 258 9413-8
14. De Meyts Bull. Mem. Acad. R Med. Belg. (1994) 149 181-90
15. Schaffer Eur. J. Biochem. (1994) 221 1127-32
16. Harmon *et al.* J. Biol. Chem. (1983) 258 6875-6881
17. Kohanski and Lane J. Biol. Chem. (1985) 260 5014-5025
18. Maturo and Hollenberg Proc. Natl. Acad. Sci. USA (1978) 75 3070-3074
19. Mortensen *et al.* Biochem. J. (1992) 281 735-743
20. Varma *et al.* Biochem Mol Biol Int (1994) 32 807-17
21. Li, et al. (1991). J. Biol. Chem. 266, 7051-7057

22. Konstantopoulos Doctor of Philosophy Thesis, Department of Medicine, University of Melbourne (1997) 288 pages
23. Isakoff *et al.* Proc. Natl. Acad. Sci. USA (1995) 92 10247-51
24. Soos *et al.* Proc. Natl. Acad. Sci. U S A (1989) 86 5217-21
- 5 25. Steele-Perkins and Roth J. Biol. Chem. (1990) 265 9458-63
26. Hem, *et al.* P Lab Animals (1998) 32(4) 364-368
27. Djuric *et al.* J. Med. Chem. (1989), 32, 1145-1147
28. Harper *et al.* J. Med. Chem. (1994), 37, 2411-2420 at 2419
29. Sawyer *et al.* J. Med. Chem. (1993), 36, 3982 - 3984, on page 3982

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS

- 1 A method for treating a patient suffering from one or more insulin related ailments, which method comprises the step of: administering to a patient a therapeutically effective amount of a compound that is a biological modulator
5 of insulin activity, which compound possesses ionic and hydrophobic chemical moieties spatially located so as to mimic the spatial location of at least an ionic and hydrophobic amino acid residue of insulin, which amino acids are associated with the binding of insulin to its receptor.
- 2 The use of at least a non-peptidyl compound, which is at least a biological
10 modulator of insulin activity and possesses ionic and hydrophobic chemical moieties spatially located so as to mimic the spatial location at least an ionic and hydrophobic amino acid residue of insulin, which amino acids are associated with the binding of insulin to its receptor; in the preparation of a medicament for the treatment of a patient suffering from one or more insulin
15 related ailments.
- 3 A method according to claim 1 or the use of at least a non-peptidyl compound according to claim 2, wherein the ionic amino acid residue is selected from the group comprising: A21 Asn, B21 Glu and A17 Glu.
- 4 A method according to claim 1 or the use of at least a non-peptidyl
20 compound according to claim 2, wherein the ionic and hydrophobic amino acid residue(s) is(are) selected from the group consisting of: A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16 Tyr, A2 Ile, A3 Val and A1 Gly.
- 5 A method according to claim 1 or the use of at least a non-peptidyl
25 compound according to claim 2, wherein at least one amino acid is selected from the group comprising: A17 Glu, B21 Glu and A21 Asn; and at least one amino acid is selected from the group comprising: B24 Phe, B25 Phe, A19 Tyr, B12 Val and B12 Tyr.

- 6 A method according to claim 1 or the use of at least a non-peptidyl compound according to claim 2, wherein the ionic and hydrophobic chemical moieties are spatially located to mimic at least one of the following groups of amino acid residues:
- 5 (i.) A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe;
(ii.) A21 Asn, B21 Glu, B24 Phe, B25 Phe;
(iii.) A21 Asn, B21 Glu, B24 Phe, B25 Phe, A1 Gly, A2 Ile, A3 Val;
(iv.) A21 Asn, B21 Glu, A17 Glu, A19 Tyr, A1 Gly, A2 Ile, A3 Val;
(v.) A21 Asn, B21 Glu, A17 Glu, B12 Val, A1 Gly, A2 Ile, A3 Val;
10 (vi.) A21 Asn, B21 Glu, B12 Val, A1 Gly, A2 Ile, A3 Val;
(vii.) A21 Asn, B21 Glu, A17 Glu, B16 Tyr, A1 Gly, A2 Ile, A3 Val;
(viii.) A21 Asn, B21 Glu, A17 Glu, A19 Tyr, B12 Val, B16 Tyr;
(ix.) A21 Asn, B21 Glu, A19 Tyr, B12 Val, B16 Tyr;
(x.) A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val,
15 B16 Tyr;
(xi.) A21 Asn, B21 Glu, B24 Phe, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
(xii.) A21 Asn, B21 Glu, B24 Phe, B25 Phe, B12 Val, B16 Tyr;
(xiii.) A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;
(xiv.) A21 Asn, B21 Glu, B24 Phe, B25 Phe, A19 Tyr;
20 (xv.) A21 Asn, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;
(xvi.) B21 Glu, A17 Glu, B24 Phe, B25 Phe, A19 Tyr;
(xvii.) A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, B12 Val;
(xviii.) A21 Asn, B21 Glu, B24 Phe, B25 Phe, B12 Val;
(xix.) A21 Asn, A17 Glu, B24 Phe, B25 Phe, B12 Val;
25 (xx.) B21 Glu, A17 Glu, B24 Phe, B25 Phe, B12 Val;
(xxi.) A21 Asn, B21 Glu, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
(xxii.) A21 Asn, B21 Glu, B24 Phe, B25 Phe, B16 Tyr;

- (xxiii.) A21 Asn, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
(xxiv.) B21 Glu, A17 Glu, B24 Phe, B25 Phe, B16 Tyr;
(xxv.) A21 Asn, B21 Glu, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
(xxvi.) A21 Asn, B21 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
5 (xxvii.) A21 Asn, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
(xxviii.) B21 Glu, A17 Glu, B24 Phe, A19 Tyr, B12 Val, B16 Tyr;
(xxix.) A21 Asn, B21 Glu, A17 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
(xxx.) A21 Asn, B21 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr;
(xxxi.) A21 Asn, A17 Glu, B25 Phe, A19 Tyr, B12 Val, B16 Tyr; or
10 (xxxii.) B21 Glu, A17 Glu, B25 Phe, A19 Tyr, B12 Val.
- 7 A method according to claim 1 or the use of at least a non-peptidyl compound according to claim 2, wherein the non-peptidyl compound has the following formula:
- $AXYXZ_n$ (formula 1)
- 15 where A is W or VXW;
V is V_1 or V_2 ;
V is substituted with up to two X groups;
 V_1 is a phenyl or 6 membered heteroaromatic ring, optionally substituted with up to 5 R_1 groups, for example but not being limited to benzene,
20 pyridine, pyridazine, pyrimidine, pyrazine, triazine;
 V_2 is a 5 member ring system which may incorporate up to 4 hetero atoms which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, for example but not being limited to cyclopenta-1,3-diene, pyrrole, furan, thiophene, oxazole, isoxazole,
25 pyrazole, imidazole, thiazole, isothiazole or triazole, the ring system being optionally substituted with up to 4 R_1 groups;
W is W_1 or W_2 or W_3 ;
W is substituted with up to two X groups;
 W_1 is V_1 ;

W₂ is a fused bicyclic ring system comprising rings of 5 or 6 atoms, which may incorporate up to 4 hetero atoms, which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R₂, oxygen or sulfur, the system being optionally substituted with up to seven R₁ groups and examples include, but are not limited to naphthalene, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, indole, benzothiophene, benzofuran, benzimidazole, indazole, benzoxazole, benzisooxazole, benzthiazole, benzisothiazole, purine, indoline, isoindoline;

W₃ is -N(R₂)R'₂;

R₁ is independently H, OH, alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy, haloalkyl, haloalkoxy, halogen, SH, thioalkyl, cyano (-CN), N(R₂)R'₂, phenyl, phenyl optionally substituted with up to five alkyl groups of 1 to 3 carbon atoms or up to five halogen atoms, benzyl, phenethyl, nitro, -COR₃, -R₅COR₃, -R₅SOR₃, -R₅SO₂R₃, -SO₂N(R₂)R'₂ or azido;

R₂ and R'₂ are independently H, alkyl of 1 to 6 carbon atoms, alkenyl of 3 to 6 carbon atoms, alkynyl of 3 to 6 carbons, hydroxyalkyl of 2 to 6 carbons, alkoxy of 2 to 6 carbons, haloalkyl, haloalkenyl, haloalkoxy, benzyl, benzyl optionally substituted with up to four R₁ groups, phenylethyl, phenylethyl optionally substituted with up to four R₁ groups, arylalkyl, and where R₂ and R'₂ can also be joined to form cyclic structures including, but not limited to pyrrolidine, piperidine, hexahydro-1H-azepine, morpholine or piperazine;

R₃ is independently H, OH, alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy, -R₄N(R₂)R'₂, mesyl, trifluoromethyl, -NHSO₂CH₃ or -NHSO₂CF₃;

R₄ is independently a bond, alkyl, alkenyl or alkynyl;

X is independently, a bond, -R₄N(R₂)R₄-, -R₄N=NR₄-, -R₄N(R₂)-N(R₂)R₄-, -R₄OR₄-, -R₄SR₄-, -R₅-, -R₅O-, -R₅S-, -R₅N(R₂)-, -SO-, sulfonyl (-SO₂-), -CO-, -CONH-, -NHCONH-, -NHCO-, -CONHCO-, -CON(R₂)-, -R₅COR₅-, -R₅COR₅N(R₂)R₅-, -N(R₂)CO- or -R₄N(R₂)R₄COR₄-;

R_5 is independently alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy;

Y is either Y_1 , Y_2 or Y_3 ;

Y is substituted with at least two, but optionally up to four X linking groups;

5 Y_1 is a fused bicyclic ring system comprising rings of 5 or 6 atoms which may incorporate up to 4 hetero atoms, which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, the ring system optionally independently incorporating a sulfoxide (SO), sulfone (SO_2) or carbonyl (CO) group and optionally up to seven R_1 groups, for example but not limited to croman, isochroman, benzofuran, cromene, 1,2,3,4-tetrahydronaphthalene, 1,4-dihydronaphthalene, indan, indene, benzopiperidine, indoline, isoindoline, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline or pteridine, coumarin or 2,3-dihydrocoumarin;

10 Y_2 is a 6:6:6 or a 6:5:6 fused tricyclic system which may incorporate up to 4 hetero atoms which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, the ring system optionally independently incorporating a sulfoxide (SO), sulfone (SO_2) or carbonyl (CO) group, and the ring system being substituted with at least two, but optionally up to four X linking groups and optionally up to seven R_1 groups and thus examples include, but are not limited to 9H-xanthone, 9H-xanthene, phenoxathiin, phenoxathiin-10-oxide, phenoxathiin-10-dioxide, acridine, phenazine, phenothiazine, phenoxazine, phenothiazine-5-oxide, phenothiazine-5-dioxide, thiathrene-5-dioxide, thiathrene-5-oxide, carbazole, dibenzo[b,d]furan, dibenzo[b,d]thiophene;


25 Y_3 is V_1 ;

Z is independently $-R_6COOH$, $-R_6SO_3H$, $-R_6NO_2$, $-R_6SO_2H$, $-R_6SO_2NHR_2$, $-R_7SO_2NHCOR_4$, -N-trifluoromesylsulfonamidate, -OH, -2-yl-

hydroxyethanoic acid ($-CH(OH)COOH$), -3-yl-2-hydroxypropanoic acid ($-CH_2CH(OH)COOH$) -2-yl-2-hydroxypropanoic acid ($-CH(CH_3)(OH)COOH$), -3-yl-2,3-dihydroxypropanoic acid ($-CH(OH)CH(OH)COOH$), -2-yl-2,3-dihydroxypropanoic acid ($-C(CH_2(OH))(OH)COOH$), -3-yl-2-hydroxypropan-3-one-1-oic acid (-

30

- COCH(OH)COOH, 2-yl-2-hydroxypropandioic acid (-
 C(COOH)(OH)COOH), -2-yl-propandioic acid (-C(COOH)(H)COOH), -4-
 yl-2-hydroxybutan-4-one-1-oic acid (-COCH₂CH(OH)COOH, 2-yl-2-
 hydroxybutan-1,4-dioic acid (-C(OH)(COOH)CH₂COOH), 3-yl-2-
 5 hydroxybutan-1,4-dioic acid (-CH(CH(OH)COOH)COOH), 5-yl-tetrazole,



- R_6 is independently a bond, alkyl, alkenyl, alkynyl, alkoxy, -CO(CH₂)_n-,
 where n is an integer between 0 and 4, alkanoic, alkenoic or alkynoic;
 with the exception that where W_1 is an optionally substituted phenyl then
 10 Y_1 cannot be an optionally substituted phenyl.

- 8 A method according to claim 7 or the use of at least a non-peptidyl
 compound according to claim 7, wherein the non-peptidyl compound is a
 dimer or heterodimer wherein the compounds are joined through a X linking
 group by way of their V or W groups.
- 15 9 A method according to claim 7 or the use of at least a non-peptidyl
 compound according to claim 7, wherein when V is V_1 or V_2 , then:
- V_1 is selected from the group consisting of, benzene, pyridine, pyridazine,
 pyrimidine, pyrazine or triazine and is optionally substituted with up to 5
 R_1 groups; and
- 20 V_2 is selected from the group consisting of, cyclopenta-1,3-diene, pyrrole,
 furan, thiophene, oxazole, isoxazole, pyrazole, imidazole, thiazole,
 isothiazole or triazole and is optionally substituted with up to 4 R_1 groups;
 and W is W_2 then
- W_2 is selected from the group consisting of naphthalene, quinoline,
 25 isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline,
 cinnoline, pteridine, indole, benzothiophene, benzofuran, benzimidazole,
 indazole, benzoxazole, benzisooxazole, benzthiazole, benzisothiazole,
 purine, indoline or isoindoline and is optionally substituted with up to
 seven R_1 groups;

and Y is either Y₁ or Y₂ then

Y₁ is selected from the group consisting of croman, isochroman, benzofuran, cromene, 1,2,3,4-tetrahydronaphthalene, 1,4-dihydronaphthalene, indan, indene, benzopiperidine, indoline, isoindoline, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline or pteridine, coumarin or 2,3-dihydrocoumarin and is optionally substituted with up to seven R₁ groups; and

Y₂ is selected from the group consisting of 9H-xanthone, 9H-xanthene, phenoxathiin, phenoxathiin-10-oxide, phenoxathiin-10-dioxide, acridine, phenazine, phenothiazine, phenoxazine, phenothiazine-5-oxide, phenothiazine-5-dioxide, thiathrene-5-dioxide, thiathrene-5-oxide, carbazole, dibenzo[b,d]furan or dibenzo[b,d]thiophene and is optionally substituted with up to seven R₁ groups.

10 A method according to claim 7 or the use of at least a non-peptidyl compound according to claim 7, wherein when A is VXW then:

V is phenyl or pyrazole, optionally substituted with up to 5 R₁ groups;

and when A is W or VXW then W is W₁, W₂ or W₃ wherein

W₁ is phenyl optionally substituted with up to 5 R₁ groups;

W₂ is naphthalene or quinoline optionally substituted with up to seven R₁ groups wherein R₁ is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;

W₃ is -N(R₂)R₂ wherein R₂ is propyl;

X is independently, a bond, methoxy (-OCH₂-), oxypropoxy (-O(CH₂)₃O-), hexenyloxy (-O(CH₂)₄CH=CH-), sulfonyloxy (-SO₂O-), methyl (-CH₂-), amidyl (-CONH-) or -NHCONH-;

and Y is either Y₁ or Y₂ then

Y₁ is croman, 4-H-chromen-4-one or naphthalene optionally substituted with up to seven R₁ groups wherein R₁ is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;

Y_2 is 9H-xanthone optionally substituted with up to seven R_1 groups wherein R_1 is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;

Y_3 is phenyl optionally substituted with up to 5 R_1 groups wherein R_1 is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl; and

Z is independently $-R_6COOH$, $-R_6SO_3H$ or $-N$ -trifluoromesylsulfonamidate wherein R_6 is independently a bond or propyl.

11 A method according to claim 7 or the use of at least a non-peptidyl
10 compound according to claim 7, wherein the non-peptidyl compound is selected from the following group of compounds:

- (i) 4,4'-Methylenebis[3-hydroxy-2-naphthalenecarboxylic acid];
- (ii) 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid;
- 15 (iii) 2,4-dichloro-6-(*N*-(trifluoromethanesulfonyl)sulfamoylphenyl 3,5-dichloro-2-hydroxybenzenesulfonate;
- (iv) 7-[(4-acetyl-3-hydroxy-2-propylphenyl)methoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid;
- (v) 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid;
- 20 (vi) 3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-(1H-pyrazol-3-yl)phenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid;
- (vii) 3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-ethoxyphenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid;
- 25 (viii) 3-[4-[7-carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]-9H-xanthene]]propanoic acid;
- (ix) 8-propyl-7-(quinol-2'-ylmethoxy)-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid;
- 30 (x) 7-(naphth-2'-ylmethoxy)-8-propyl-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid;

- (xi) *N*-(trifluoromethanesulfonyl)-3,5-dinitro-4-(*N,N*-dipropylamino)benzenesulfonamide;
- (xii) 8-propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid;
- (xiii) 3,4-dihydro-7-[[6-(4-methoxyphenyl)hexenyl]oxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid; or
- (xiv) 8,8'-[Carbonylbis[imino-3,1-phenylenecarbonylimino(4-methyl-3,1-phenylene)carbonylimino]]bis-1,3,5-naphthalenetrisulfonic acid.
- 10 12 A pharmaceutical composition comprising at least a chemical compound capable of modulating the biological activity of insulin and a pharmaceutically acceptable carrier and/or diluent; wherein said compound has the following general formula.
- $$\text{AXYXZ}_n \quad (\text{formula 1})$$
- 15 where A is W or VXW;
V is V_1 or V_2 ;
V is substituted with up to two X groups;
 V_1 is a phenyl or 6 membered heteroaromatic ring, optionally substituted with up to 5 R_1 groups, for example but not being limited to benzene,
20 pyridine, pyridazine, pyrimidine, pyrazine, triazine;
 V_2 is a 5 member ring system which may incorporate up to 4 hetero atoms which may be independently a nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, for example but not being limited to cyclopenta-1,3-diene, pyrrole, furan, thiophene, oxazole, isoxazole,
25 pyrazole, imidazole, thiazole, isothiazole or triazole, the ring system being optionally substituted with up to 4 R_1 groups;
W is W_1 or W_2 or W_3 ;
W is substituted with up to two X groups;
 W_1 is V_1 ;
30 W_2 is a fused bicyclic ring system comprising rings of 5 or 6 atoms, which may incorporate up to 4 hetero atoms, which may be independently a

nitrogen atom, a nitrogen atom optionally substituted with R_2 , oxygen or sulfur, the system being optionally substituted with up to seven R_1 groups and examples include, but are not limited to naphthalene, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, indole, benzothiophene, benzofuran, benzimidazole, indazole, benzoxazole, benzisooxazole, benzthiazole, benzisothiazole, purine, indoline, isoindoline;

W_3 is $-N(R_2)R'_2$;

R_1 is independently H, OH, alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy, haloalkyl, haloalkoxy, halogen, SH, thioalkyl, cyano ($-CN$), $N(R_2)R'_2$, phenyl, phenyl optionally substituted with up to five alkyl groups of 1 to 3 carbon atoms or up to five halogen atoms, benzyl, phenethyl, nitro, $-COR_3$, $-R_5COR_3$, $-R_5SOR_3$, $-R_5SO_2R_3$, $-SO_2N(R_2)R'_2$ or azido;

R_2 and R'_2 are independently H, alkyl of 1 to 6 carbon atoms, alkenyl of 3 to 6 carbon atoms, alkynyl of 3 to 6 carbons, hydroxyalkyl of 2 to 6 carbons, alkoxy of 2 to 6 carbons, haloalkyl, haloalkenyl, haloalkoxy, benzyl, benzyl optionally substituted with up to four R_1 groups, phenylethyl, phenylethyl optionally substituted with up to four R_1 groups, arylalkyl, and where R_2 and R'_2 can also be joined to form cyclic structures including, but not limited to pyrrolidine, piperidine, hexahydro-1H-azepine, morpholine or piperazine;

R_3 is independently H, OH, alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy, $-R_4N(R_2)R'_2$, mesyl, trifluoromethyl, $-NHSO_2CH_3$ or $-NHSO_2CF_3$;

R_4 is independently a bond, alkyl, alkenyl or alkynyl;


X is independently, a bond, $-R_4N(R_2)R_4-$, $-R_4N=NR_4-$, $-R_4N(R_2)-N(R_2)R_4-$, $-R_4OR_4-$, $-R_4SR_4-$, $-R_5-$, $-R_5O-$, $-R_5S-$, $-R_5N(R_2)-$, $-SO-$, sulfonyl ($-SO_2-$), $-CO-$, $-CONH-$, $-NHCONH-$, $-NHCO-$, $-CONHCO-$, $-CON(R_2)-$, $-R_5COR_5-$, $-R_5COR_5N(R_2)R_5-$, $-N(R_2)CO-$ or $-R_4N(R_2)R_4COR_4-$;

R_5 is independently alkyl, alkenyl, alkynyl, alkoxy, alkanol, hydroxyalkoxy;

Y is either Y_1 , Y_2 or Y_3 ;

Y is substituted with at least two, but optionally up to four X linking groups;
Y₁ is a fused bicyclic ring system comprising rings of 5 or 6 atoms which
may incorporate up to 4 hetero atoms, which may be independently a
nitrogen atom, a nitrogen atom optionally substituted with R₂, oxygen or
sulfur, the ring system optionally independently incorporating a sulfoxide
(SO), sulfone (SO₂) or carbonyl (CO) group and optionally up to seven R₁
groups, for example but not limited to croman, isochroman, benzofuran,
cromene, 1,2,3,4-tetrahydronaphthalene, 1,4-dihydronaphthalene, indan,
indene, benzopiperidine, indoline, isoindoline, quinoline, isoquinoline,
phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline or
pteridine, coumarin or 2,3-dihydrocoumarin;
Y₂ is a 6:6:6 or a 6:5:6 fused tricyclic system which may incorporate up to
4 hetero atoms which may be independently a nitrogen atom, a nitrogen
atom optionally substituted with R₂, oxygen or sulfur, the ring system
optionally independently incorporating a sulfoxide (SO), sulfone (SO₂) or
carbonyl (CO) group, and the ring system being substituted with at least
two, but optionally up to four X linking groups and optionally up to seven
R₁ groups and thus examples include, but are not limited to 9H-xanthone,
9H-xanthene, phenoxathiin, phenoxathiin-10-oxide, phenoxathiin-10-
dioxide, acridine, phenazine, phenothiazine, phenoxazine, phenothiazine-
5-oxide, phenothiazine-5-dioxide, thiathrene-5-dioxide, thiathrene-5-oxide,
carbazole, dibenzo[b,d]furan, dibenzo[b,d]thiophene;
Y₃ is V₁;
Z is independently -R₆COOH, -R₆SO₃H, -R₆NO₂, -R₆SO₂H, -R₆SO₂NHR₂;
-R₇SO₂NHCOR₄, -N-trifluoromesylsulfonamidate, -OH, -2-yl-
hydroxyethanoic acid (-CH(OH)COOH), -3-yl-2-hydroxypropanoic acid (-
CH₂CH(OH)COOH) -2-yl-2-hydroxypropanoic acid (-
CH(CH₃)(OH)COOH), -3-yl-2,3-dihydroxypropanoic acid (-
CH(OH)CH(OH)COOH), -2-yl-2,3-dihydroxypropanoic acid (-
C(CH₂(OH))(OH)COOH), -3-yl-2-hydroxypropan-3-one-1-oic acid (-
COCH(OH)COOH, 2-yl-2-hydroxypropandioic acid (-
C(COOH)(OH)COOH), -2-yl-propandioic acid (-C(COOH)(H)COOH), -4-

yl-2-hydroxybutan-4-one-1-oic acid ($-\text{COCH}_2\text{CH}(\text{OH})\text{COOH}$), 2-yl-2-hydroxybutan-1,4-dioic acid ($-\text{C}(\text{OH})(\text{COOH})\text{CH}_2\text{COOH}$), 3-yl-2-hydroxybutan-1,4-dioic acid ($-\text{CH}(\text{CH}(\text{OH})\text{COOH})\text{COOH}$), 5-yl-tetrazole,



- 5 R_6 is independently a bond, alkyl, alkenyl, alkynyl, alkoxy, $-\text{CO}(\text{CH}_2)_n-$, where n is an integer between 0 and 4, alkanoic, alkenoic or alkynoic; with the exception that where W_1 is an optionally substituted phenyl then Y_1 cannot be an optionally substituted phenyl.

- 10 13 A pharmaceutical composition according to claim 12, wherein the non-peptidyl compound is a dimers or heterodimers of compounds where such compounds are joined through a X linking group by way of their V or W groups.

- 14 A pharmaceutical composition according to claim 12, wherein when V is V_1 or V_2 , then:

- 15 V_1 is selected from the group consisting of, benzene, pyridine, pyridazine, pyrimidine, pyrazine or triazine and is optionally substituted with up to 5 R_1 groups; and

- 20 V_2 is selected from the group consisting of, cyclopenta-1,3-diene, pyrrole, furan, thiophene, oxazole, isoxazole, pyrazole, imidazole, thiazole, isothiazole or triazole and is optionally substituted with up to 4 R_1 groups;

and W is W_2 then

- 25 W_2 is selected from the group consisting of naphthalene, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, indole, benzothiophene, benzofuran, benzimidazole, indazole, benzoxazole, benzisooxazole, benzthiazole, benzisothiazole, purine, indoline or isoindoline and is optionally substituted with up to seven R_1 groups;

and Y is either Y_1 or Y_2 then

Y₁ is selected from the group consisting of croman, isochroman, benzofuran, cromene, 1,2,3,4-tetrahydronaphthalene, 1,4-dihydronaphthalene, indan, indene, benzopiperidine, indoline, isoindoline, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline or pteridine, coumarin or 2,3-dihydrocoumarin and is optionally substituted with up to seven R₁ groups; and

Y₂ is selected from the group consisting of 9H-xanthone, 9H-xanthene, phenoxathiin, phenoxathiin-10-oxide, phenoxathiin-10-dioxide, acridine, phenazine, phenothiazine, phenoxazine, phenothiazine-5-oxide, phenothiazine-5-dioxide, thiathrene-5-dioxide, thiathrene-5-oxide, carbazole, dibenzo[b,d]furan or dibenzo[b,d]thiophene and is optionally substituted with up to seven R₁ groups.

15 A pharmaceutical composition according to claim 12, wherein in the non-peptidyl compound of formula 1 when A is W or VXW then:

15 V is phenyl or pyrazole, optionally substituted with up to 5 R₁ groups; and W is W₁, W₂ or W₃ then

W₁ is phenyl optionally substituted with up to 5 R₁ groups;

W₂ is naphthalene or quinoline optionally substituted with up to seven R₁ groups wherein R₁ is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;

W₃ is -N(R₂)R₂ wherein R₂ is propyl;

X is independently, a bond, methoxy (-OCH₂-), oxypropoxy (-O(CH₂)₃O-), hexenyloxy (-O(CH₂)₄CH=CH-), sulfonyloxy (-SO₂O-), methyl (-CH₂-), amidyl (-CONH-) or -NHCONH-;

25 and Y is either Y₁ or Y₂ then

Y₁ is croman, 4-H-chromen-4-one or naphthalene optionally substituted with up to seven R₁ groups wherein R₁ is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;

Y_2 is 9H-xanthone optionally substituted with up to seven R_1 groups wherein R_1 is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl;

5 Y_3 is phenyl optionally substituted with up to 5 R_1 groups wherein R_1 is independently H, OH, methyl, ethyl, propyl, nitro, methoxy, ethoxy, 2-hydroxyethoxy, chloro, fluoro or acetyl; and

Z is independently $-R_6COOH$, $-R_6SO_3H$ or $-N$ -trifluoromesylsulfonamidate wherein R_6 is independently a bond or propyl.

16 A pharmaceutical composition according to claim 12, wherein the non-peptidyl compound is selected from the following group of compounds:

- (xv) 4,4'-Methylenebis[3-hydroxy-2-naphthalenecarboxylic acid];
- (xvi) 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid;
- (xvii) 2,4-dichloro-6-(*N*-(trifluoromethanesulfonyl))sulfamoylphenyl 3,5-dichloro-2-hydroxybenzenesulfonate;
- 15 (xviii) 7-[(4-acetyl-3-hydroxy-2-propylphenyl)methoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid;
- (xix) 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid;
- 20 (xx) 3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-(1H-pyrazol-3-yl)phenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid;
- (xxi) 3,4-dihydro-8-propyl-7-[[3-[2-ethyl-5-hydroxy-4-ethoxyphenoxy]propyl]oxy]-2H-1-benzopyran-2-carboxylic acid;
- (xxii) 3-[4-[7-carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]-9H-xanthene]]propanoic acid;
- 25 (xxiii) 8-propyl-7-(quinol-2'-ylmethoxy)-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid;
- (xxiv) 7-(naphth-2'-ylmethoxy)-8-propyl-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid;

- (xxv) *N*-(trifluoromethanesulfonyl)-3,5-dinitro-4-(*N,N'*-dipropylamino)benzenesulfonamide;
- (xxvi) 8-propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid;
- (xxvii) 3,4-dihydro-7-[[6-(4-methoxyphenyl)hexenyl]oxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid; or
- (xxviii) 8,8'-[Carbonylbis[imino-3,1-phenylenecarbonylimino(4-methyl-3,1-phenylene)carbonylimino]]bis-1,3,5-naphthalenetrisulfonic acid.
- 10 17 A method for identifying a non-peptidyl compound possessing ionic and hydrophobic chemical moieties spatially located so as to mimic particular ionic and hydrophobic amino acid residues of insulin which are associated with the binding of insulin to its receptor, said method comprising the steps of: (1) comparing the three dimensional structure of the non-peptidyl compound with a three dimensional pharmacophore of an active site of insulin; and (2) selecting a non-peptidyl compound with ionic and hydrophobic chemical moieties spatially located so as to mimic said site
- 15
- 18 A method for determining whether a non-peptidyl compound identified according to the method of claim 17 is an agonist or an antagonist, said method comprising the step of: exposing the compound to an insulin or insulin like receptor and measuring the change in biological activity following exposure of the compound to the receptor.
- 20
- 19 A method according to claim 1 as substantially herein before described.
- 20
- 20 The use of a compound according to claim 2 as substantially herein before described.
- 25
- 21 A pharmaceutical composition according to claim 12 as substantially herein before described.

| Hypothesis | Residue | | | | | | | | | | |
|------------|---------|-----|-----|-----|-----|-----|-----|-----|----|----|----|
| | A21 | B21 | A17 | B24 | B25 | A19 | B12 | B16 | A1 | A2 | A3 |
| 1 | | | | | | | | | | | |
| 2 (1a) | | | | | | | | | | | |
| 3 | | | | | | | | | | | |
| 4 | | | | | | | | | | | |
| 5 | | | | | | | | | | | |
| 5a | | | | | | | | | | | |
| 6 | | | | | | | | | | | |
| 7 | | | | | | | | | | | |
| 7a | | | | | | | | | | | |
| 8 | | | | | | | | | | | |
| 8a | | | | | | | | | | | |
| 8d | | | | | | | | | | | |
| 9 | | | | | | | | | | | |
| 9a | | | | | | | | | | | |
| 9b | | | | | | | | | | | |
| 9e | | | | | | | | | | | |
| 10 | | | | | | | | | | | |
| 10a | | | | | | | | | | | |
| 10b | | | | | | | | | | | |
| 10e | | | | | | | | | | | |
| 11 | | | | | | | | | | | |
| 11a | | | | | | | | | | | |
| 11b | | | | | | | | | | | |
| 11e | | | | | | | | | | | |
| 12 | | | | | | | | | | | |
| 12a | | | | | | | | | | | |
| 12b | | | | | | | | | | | |
| 12e | | | | | | | | | | | |
| 13 | | | | | | | | | | | |
| 13a | | | | | | | | | | | |
| 13b | | | | | | | | | | | |
| 13e | | | | | | | | | | | |

Figure 2

| Residue | Feature | Tolerance HIU A | Tolerance HIT3 A | Tolerance HIT4 A |
|----------------------------------|-----------------------|--------------------|---------------------|---------------------|
| A17 COO ⁻ | Negative ionisable | 5 | 3 | 3 |
| A21 COO ⁻ | Negative ionisable | 5 | 3 | 3 |
| B21 COO ⁻ | Negative ionisable | 5 | 3 | 3 |
| B24 Phe | Hydrophobic | 5 | 3 | 3 |
| B25 Phe | Hydrophobic | 5 | 3 | 3 |
| A 1 NH ₃ ⁺ | Positive ionisable | 5 | 3 | 3 |
| A2 Ile | Hydrophobic | 5 | 3 | 3 |
| A3 Val | Hydrophobic | 3 | 3 | 3 |
| A19 Tyr | Hydrophobic | 3.6 | 3 | 3 |
| B16 Tyr | Hydrophobic | 3.6 | 3 | 3 |
| B12 Val | Hydrophobic | 3.6 | 3 | 3 |

Figure 3

| Hypothesis | Tolerance | | | |
|------------|-----------|----|----|----|
| | 1.5A | 2A | 3A | 5A |
| 1 | | | | |
| 2 (1a) | | | | |
| 3 | | | | |
| 4 | | | | |
| 5 | | | | |
| 5a | | | | |
| 6 | | | | |
| 7 | | | | |
| 7a | | | | |
| 8 | | | | |
| 8a | | | | |
| 8d | | | | |
| 9 | | | | |
| 9a | | | | |
| 9b | | | | |
| 9e | | | | |
| 10 | | | | |
| 10a | | | | |
| 10b | | | | |
| 10e | | | | |
| 11 | | | | |
| 11a | | | | |
| 11b | | | | |
| 11e | | | | |
| 12 | | | | |
| 12a | | | | |
| 12b | | | | |
| 12e | | | | |
| 13 | | | | |
| 13a | | | | |
| 13b | | | | |
| 13e | | | | |

Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12

9/35




Figure 13




Figure 14

10/35




Figure 15




Figure 16

11/35




Figure 17




Figure 18

12/35




Figure 19




Figure 20

13/35




Figure 21




Figure 22

14/35




Figure 23




Figure 24

15/35




Figure 25




Figure 26

16/35




Figure 27




Figure 28

17/35




Figure 29




Figure 30

18/35




Figure 31




Figure 32

19/35




Figure 33




Figure 34

20/35




Figure 35




Figure 36

21/35




Figure 37




Figure 38

22/35




Figure 39




Figure 40

23/35




Figure 41




Figure 42

24/35




Figure 43




Figure 44

25/35




Figure 45




Figure 46

26/35




Figure 47




Figure 48




Figure 49




Figure 50




Figure 51




Figure 52

29/35




Figure 53




Figure 54

30/35




Figure 55




Figure 56

31/35




Figure 57




Figure 58

32/35





Figure 59

33/35




Figure 60




Figure 61

34/35




Figure 62




Figure 63

35/35




Figure 64




Figure 65

International application No.
PCT/AU 99/00786

Int Cl⁶: A61K 38/28, 31/19, 31/35

B. FIELDS SEARCHED

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
AU: IPC as above

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| X | AU 86/54495 (612964) (NOVO NORDISK A/S) 18 September 1986 | 1-6,17,18 |
| X | AU 86/62066 (593274) (NOVO NORDISK A/S) 3 March 1987 | 1-6,17,18 |
| X | AU 88/13976 (612141) (NOVO NORDISK A/S) 26 September 1988 | 1-6,17,18 |

☒ Further documents are listed in the continuation of Box C

☒ See patent family annex

| | |
|-----|---|
| "A" | Document defining the general state of the art which is not considered to be of particular relevance |
| "E" | earlier application or patent but published on or after the international filing date |
| "L" | document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) |
| "O" | document referring to an oral disclosure, use, exhibition or other means |
| "P" | document published prior to the international filing date but later than the priority date claimed |

| | |
|-----|--|
| "T" | later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| "X" | document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| "Y" | document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| "&" | document member of the same patent family |

Date of the actual completion of the international search
10 December 1999

Date of mailing of the international search report
01 FEB 2000

Name and mailing address of the ISA/AU
AUSTRALIAN PATENT OFFICE
PO BOX 200
WODEN ACT 2606 AUSTRALIA
E-mail address: pct@ipaustralia.gov.au
Facsimile No.: (02) 6285 3929

Authorized officer

A. WILCOX
Telephone No.: (02) 6283 2243

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU 99/00786

| C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT | | |
|---|---|-----------------------|
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| X | AU 88/22636 (616205) (NOVO NORDISK A/S) 19 February 1990 | 1-6,17,18 |
| X | AU 90/48344 (641631) (NOVO NORDISK A/S) 1 August 1990 | 1-6,17,18 |
| X | AU 90/55415 (631868) (MOUNT SINAI SCHOOL OF MEDICINE) 16 November 1990 | 1-6,17,18 |
| X | WO 95/17183 (ELI LILLEY AND COMPANY) 29 June 1995 | 1,12-16,21 |
| X | EP 132366 (ELI LILLEY AND COMPANY) 30 February 1985 | 12-16,21 |
| X | SAWYER, J S et al: Biphenyl-Substituted Xanthenes: Highly Potent Leukotriene B ₄ Receptors Antagonists. J. Med. Chem. 1993, volume 36: 3982-3984 | 12-16,21 |
| X | HARPER, R W et al: Leukotriene B ₄ (LTB ₄) Receptor Antagonists: A series of (Hydroxyphenylpyrazoles). J. Med. Chem. 1994 volume 37: 2411-2420 See page 2419, paragraph 4, lines 28-44 | 12-16,21 |
| X | DJURIC, S W et al; 7-[3-(4-Acetyl-3-methoxy-2-propylphenoxy) propoxy]- 3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid: An Orally Active Selective Leukotriene B ₄ Receptor Antagonist. J. Med. Chem, 1989 volume 32, No: 6: 1145-47 | 12-16,21 |
| X | Chemical Abstracts, Abstract No: 490788, 1990, Aguilar-Bryan, L, et al, Photoaffinity labelling and partial purification of the beta cell sulfonylurea receptor using a novel, biologically active glyburide analog. J. Biol. Chem, 1990 volume 265, No: 14: 8218-24 | 7,9,10 |
| X | Chemical Abstracts, Abstract No: 426785 1980 Bahn H, et al, Cystic Acid as a solubilizing protective group in peptide chemistry. Schriftenr Dtsch Wolfforschungsinst 1978, 76: 5-7 | 7,9,10 |

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/AU 99/00786

Box I Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

Box II Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See Continuation of Box II.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU 99/00786

Box II (continued)

The claims do not comply with the requirements of Unity of Invention. The basic test in determining Unity of Invention is set out in rules 13.2 to 13.4. The major requirement for Unity of Invention is that there is a technical relationship among those inventions involving one or more of the same or corresponding special technical features and that the technical features are novel and have an inventive step when considered in the light of the prior art. Both a priori lack of unity (whether the claims have the same or corresponding special technical features) and a posteriori lack of unity (whether the common features are novel or involve an inventive step).

Claims 1-6, 19 and 20 are directed to a method of treatment using a biological modulator of insulin activity. The biological modulator of insulin is an insulin molecule modified at the following amino acids: A1, A2, A3, A17, A19, A20, A21, B12, B16, B21, B22 and B24. The different sequences are considered to be different technical features and, a priori, lack unity. The search disclosed that the peptide mimics of insulin and the use of such peptides as modulators of insulin activity are well known and Claims 1-6, 19 and 20, a posteriori, lack unity.

Claims 7-16 and 21 are directed to pharmaceutical formulations of compounds of Formula I and the use of such formulations as modulators of insulin activity. The search disclosed that compounds of Formula I and the use of compounds of Formula I as modulators of insulin activity are well known. Formula I includes a very large number of compounds, and the compounds are not considered to have the same or corresponding technical features. Claims 7-16 and 21 are considered to lack unity of invention.

Claims 17 and 18 are directed to a method of identifying peptide mimics of the spatial conformation of insulin. Claims 17 and 18 do not have the same special technical features as claims 1-16 and 19-21.

There is no significant structural element shared by all of the alternatives described in Claims 1-6 and in Formula I and therefore, unity of invention cannot be established.

The search report has been established as follows:

Claims 1-6, 19 and 20 have been fully searched. The claims are directed to exemplified compounds. Claims 7-16 and 21 have been partially searched based on a search of the exemplified compounds.

Information on patent family members

PCT/AU 99/00786

| Patent Document Cited in Search Report | | | | Patent Family Member | | | |
|--|----------|----|----------|----------------------|----------|----|----------|
| AU | 54495 | CN | 86101489 | CS | 8601666 | DD | 244345 |
| | | DK | 1070/86 | EP | 194864 | ES | 552879 |
| | | ES | 8800276 | FI | 861007 | GR | 860687 |
| | | HU | 41052 | IL | 78103 | JP | 61212598 |
| | | NO | 860920 | NZ | 215445 | PH | 24260 |
| | | PT | 82173 | US | 4946828 | US | 5008241 |
| | | YU | 345/86 | ZA | 8601432 | AU | 75916/87 |
| | | CN | 87104988 | DD | 273839 | DK | 3761/87 |
| | | EP | 254516 | FI | 873185 | HU | 45271 |
| | | IL | 83243 | NO | 873018 | NZ | 221130 |
| | | PT | 85355 | YU | 1353/87 | ZA | 8705295 |
| | | AU | 13976/88 | DK | 5903/88 | EP | 280534 |
| | | FI | 884914 | NO | 884724 | NZ | 223630 |
| | | PT | 86819 | WO | 88/06599 | ZA | 8801280 |
| | | DK | 5400/87 | AU | 79907/87 | CA | 1316819 |
| | | CN | 87107049 | EP | 265213 | FI | 874596 |
| | | HU | 46232 | IL | 84203 | JP | 63179831 |
| | | NO | 874347 | NZ | 222214 | PH | 23446 |
| | | PT | 85949 | US | 5070186 | YU | 1918/87 |
| | | DD | 266964 | ZA | 8707825 | | |
| AU | 86/62066 | CA | 1306212 | CN | 86106574 | DK | 4116/86 |
| | | EP | 214826 | FI | 863512 | GR | 862233 |
| | | HU | 42526 | IL | 79887 | JP | 62053999 |
| | | NO | 863474 | NZ | 217406 | PH | 25772 |
| | | PT | 83278 | US | 5618913 | YU | 41/88 |
| | | YU | 1484/86 | CS | 8606310 | DD | 268976 |
| | | ZA | 8606450 | DK | 3965/85 | EP | 175607 |
| | | ES | 546468 | ES | 8702487 | FR | 2582670 |
| | | JP | 61063277 | FR | 2569719 | | |

CONTINUED

CONTINUED

INTERNATIONAL SEARCH REPORT

| Patent Document Cited in Search Report | | | | Patent Family Member | | | |
|--|----------|----|----------|----------------------|----------|----|----------|
| AU | 88/13976 | DK | 5400/87 | DK | 5903/88 | EP | 280534 |
| | | FI | 884914 | NO | 884724 | NZ | 223630 |
| | | PT | 86819 | WO | 88/06599 | AU | 79907/87 |
| | | CA | 1316819 | CN | 87107049 | EP | 265213 |
| | | FI | 874596 | HU | 46232 | IL | 84203 |
| | | JP | 63179831 | NO | 874347 | NZ | 222214 |
| | | PH | 23446 | PT | 85949 | US | 5070186 |
| | | YU | 1918/87 | US | 5008241 | ZA | 8801280 |
| | | DD | 266964 | ZA | 8707825 | AU | 75916/87 |
| | | CN | 87104988 | DD | 273839 | DK | 3761/87 |
| | | EP | 254516 | FI | 873185 | HU | 45271 |
| | | IL | 83243 | JP | 63099096 | NO | 873018 |
| | | NZ | 221130 | PT | 85355 | US | 4946828 |
| | | YU | 1353/87 | ZA | 8705295 | DK | 1070/86 |
| | | AU | 54495/86 | CN | 86101489 | CS | 8601666 |
| | | DD | 244345 | EP | 194864 | ES | 552879 |
| | | ES | 8800276 | FI | 861007 | GR | 860687 |
| | | HU | 41052 | IL | 78103 | JP | 61212598 |
| | | NO | 860920 | NZ | 215445 | PH | 24260 |
| | | PT | 82173 | YU | 345/86 | ZA | 8601432 |
| AU | 88/22636 | WO | 90/01038 | EP | 425482 | NO | 905472 |
| | | US | 5149777 | | | | |
| AU | 90/48344 | CA | 2006578 | CN | 1043719 | DK | 1193/91 |
| | | EP | 375437 | EP | 837072 | EP | 861851 |
| | | HU | 56857 | NO | 912438 | NZ | 231936 |
| | | US | 5164366 | US | 5716927 | WO | 90/07522 |
| | | YU | 2436/89 | DD | 290426 | ZA | 8909501 |
| AU | 90/55415 | CA | 2014896 | EP | 469084 | GR | 90100290 |
| | | HU | 59941 | IL | 94163 | IL | 111437 |
| | | NO | 914085 | PT | 93832 | WO | 90/12814 |
| | | ZA | 9002965 | US | 5208217 | | |
| WO | 95/17183 | AU | 14028/95 | BR | 9408407 | CA | 2179649 |
| | | CN | 1142768 | CZ | 9601780 | EP | 735870 |
| | | FI | 962557 | HU | 75335 | NO | 962568 |
| | | US | 5478857 | ZA | 9410041 | US | 5563164 |

CONTINUED

INTERNATIONAL SEARCH REPORT

| Patent Document Cited in Search Report | | | | Patent Family Member | | | |
|--|--------|----|----------|----------------------|---------|----|---------|
| EP | 132366 | AU | 30751/84 | CA | 1242196 | CS | 8405487 |
| | | DD | 229116 | DK | 3491/84 | ES | 534268 |
| | | ES | 8601836 | FI | 842844 | GB | 2143530 |
| | | GR | 81529 | HU | 44472 | IL | 72384 |
| | | JP | 60048944 | NZ | 208880 | PT | 78921 |
| | | SU | 1516011 | US | 5105017 | ZA | 8405408 |
| | | US | 5393768 | | | | |

END OF ANNEX

THIS PAGE BLANK (USPTO)